

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 December 2001 (13.12.2001)

PCT

(10) International Publication Number  
**WO 01/94345 A2**

(51) International Patent Classification<sup>7</sup>: **C07D 471/00**

(21) International Application Number: PCT/US01/15936

(22) International Filing Date: 15 May 2001 (15.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/210,137 7 June 2000 (07.06.2000) US

(71) Applicant (for all designated States except US): **LILLY ICOS LLC** [US/US]; 1209 Orange Street, Wilmington, DE 19801 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ORME, Mark, W.** [US/US]; 4235 Francis Avenue, #203, Seattle, WA 98103 (US). **SAWYER, Jason, Scott** [US/US]; 5718 North Winthrop Avenue, Indianapolis, IN 46220 (US). **DAUGAN, Alain, Claud-Marie** [FR/FR]; Centre de Recherche Glaxo Smith Kline Courtaboeuf, ZA de Courtaboeuf, 25, avenue de Quebec, F-91940 Les Ulis (FR).

(74) Agent: **NAPOLI, James, J.**; Marshall, O'Toole, Gerstein, Murray & Borun, 6300 Sears Tower, 233 South Wacker Drive, Chicago, IL 60606 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**


— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/94345 A2

(54) Title: CHEMICAL COMPOUNDS



(57) Abstract: Compounds of the general structural formula (I) and use of the compounds and salts and solvates thereof, as therapeutic agents.

- 1 -


## CHEMICAL COMPOUNDS

FIELD AND BACKGROUND OF THE INVENTION

5 This invention relates to a series of compounds, to methods of preparing the compounds, to pharmaceutical compositions containing the compounds, and to their use as therapeutic agents. In particular, the PRESENT invention relates to compounds that are potent and selective inhibitors of cyclic guanosine 3',5'-monophosphate specific phosphodiesterase (cGMP-specific PDE), in particular PDE5, and have utility in a variety of therapeutic areas wherein such inhibition is considered beneficial, including the treatment of cardiovascular disorders and erectile dysfunction.

DETAILED DESCRIPTION OF THE INVENTION

20 The present invention is directed to compounds having the general structural formula (I):



(I)


30 wherein R<sup>1</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, haloC<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl,

- 2 -

heteroC<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkylC<sub>1-3</sub>alkyl, arylC<sub>1-3</sub>alkyl, and heteroarylC<sub>1-3</sub>alkyl;

R<sup>2</sup> is selected from the group consisting of an optionally substituted monocyclic aromatic ring  
5 selected from the group consisting of benzene, thiophene, furan, and pyridine, and an optionally substituted bicyclic ring

10



15

wherein the fused ring A is a 5- or 6-membered ring, saturated or partially or fully unsaturated, and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulfur, and nitrogen;

20

R<sup>3</sup> is selected from the group consisting of hydrogen and C<sub>1-6</sub>alkyl,

or R<sup>1</sup> and R<sup>3</sup> together represent a 3- or 4-membered alkyl or alkenyl chain component of a 5- or 6-membered ring;

25

R<sup>4</sup>, independently, is selected from the group consisting of

aryl,

Het,

C<sub>3-8</sub>cycloalkyl,

YC<sub>3-8</sub>cycloalkyl (wherein Y is oxygen, sul-

30

fur, or NR<sup>a</sup>),

C(=O)R<sup>a</sup>,

OC(=O)R<sup>a</sup>,

C(=O)OR<sup>a</sup>,

- 3 -

$C_{1-4}alkyleneNR^aR^b$ ,  
 $C_{1-4}alkyleneHet$ ,  
 $C_{1-4}alkyleneC(=O)OR^a$ ,  
 $C(=O)NR^aSO_2R^c$ ,  
5  $C(=O)C_{1-4}alkyleneHet$ ,  
 $C(=O)NR^aR^b$ ,  
 $C(=O)NR^aR^c$ ,  
 $C(=O)NR^aC_{1-4}alkyleneOR^b$ ,  
 $C(=O)NR^aC_{1-4}alkyleneHet$ ,  
10  $OR^a$ ,  
 $OC_{1-4}alkyleneC(=O)OR^a$ ,  
 $OC_{2-4}alkyleneNR^aR^b$ ,  
 $OC_{1-4}alkyleneHet$ ,  
 $OC_{2-4}alkyleneOR^a$ ,  
15  $OC_{1-4}alkyleneNR^aC(=O)OR^b$ ,  
 $NR^aR^b$ ,  
 $NR^aC_{1-4}alkyleneNR^aR^b$ ,  
 $NR^aC(=O)R^b$ ,  
 $NR^aC(=O)NR^aR^b$ ,  
20  $N(SO_2C_{1-4}alkyl)_2$ ,  
 $NR^a(SO_2C_{1-4}alkyl)$ ,  
nitro ( $NO_2$ ),  
trifluoromethyl,  
trifluoromethoxy,  
25 cyano ( $CN$ ),  
 $SO_2NR^aR^b$ ,  
 $SO_2R^a$ ,  
 $SOR^a$ ,  
 $SR^a$ ,  
30 and  $OSO_2CF_3$ ;  
 $R^5$  is selected from the group consisting of  
hydrogen, halogen, and  $C_{1-6}alkyl$ ;

- 4 -

or R<sup>4</sup> and R<sup>5</sup> are taken together with the carbon atoms to which they are attached to form a 5-, 6-, or 7-membered ring, saturated or partially or fully unsaturated, optionally substituted and optionally containing one or two heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen;

R<sup>a</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, aryl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, and Het;

R<sup>b</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkylenearyl, and Het;

R<sup>c</sup> is phenyl or C<sub>4-6</sub>cycloalkyl, either optionally substituted with one or more substituents selected from the group consisting of halo, C(=O)OR<sup>a</sup>, and OR<sup>a</sup>;

Het is a 5- or 6-membered heterocyclic group, saturated or partially or fully unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with C<sub>1-4</sub>alkyl or C(=O)OR<sup>b</sup>;

q is 1, 2, or 3; and

pharmaceutically acceptable salts and hydrates thereof.

As used herein, the term "alkyl" includes straight chained and branched hydrocarbon groups containing the indicated number of carbon atoms, typically methyl, ethyl, and straight chain and branched propyl and butyl groups. The hydrocarbon group can contain up to 16 carbon atoms. The term "alkyl" includes "bridged alkyl," i.e., a C<sub>6</sub>-C<sub>16</sub> bicyclic or polycyclic hydrocarbon group, for exam-

- 5 -

ple, norbornyl, adamantyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, bicyclo[3.2.1]octyl, or decahydronaphthyl. The term "cycloalkyl" is defined as a cyclic C<sub>3</sub>-C<sub>8</sub> hydrocarbon group, e.g., cyclopropyl, 5 cyclobutyl, cyclohexyl, and cyclopentyl.

"Heterocycloalkyl" is similarly defined as a ring containing three to eight atoms, with one to three atoms selected from the group consisting of oxygen, nitrogen, and sulfur, with the remaining 10 atoms being carbon.

The terms "alkenyl" and "alkynyl" are defined identically as "alkyl," except for containing a carbon-carbon double bond or carbon-carbon triple bond, respectively.

15 The term "alkylene" refers to an alkyl group having a substituent. For example, the term "C<sub>1-3</sub>alkylenearyl" refers to an alkyl group containing one to three carbon atoms, and substituted with an aryl group. The term "alkenylene" as used herein 20 is similarly defined, and contains the indicated number of carbon atoms and a carbon-carbon double bond, and includes straight chained and branched alkenylene groups, like ethylenylene.

The term "halo" or "halogen" is defined 25 herein to include fluorine, bromine, chlorine, and iodine.

The term "haloalkyl" is defined herein as an alkyl group substituted with one or more halo substituents, either fluoro, chloro, bromo, iodo, or 30 combinations thereof. Similarly, "halocycloalkyl" is defined as a cycloalkyl group having one or more halo substituents.

- 6 -

The term "aryl," alone or in combination, is defined herein as a monocyclic or polycyclic aromatic group, preferably a monocyclic or bicyclic aromatic group, e.g., phenyl or naphthyl, that can be unsubstituted or substituted, for example, with one or more, and in particular one to three, halo, alkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkyl, nitro, amino, alkylamino, acylamino, alkylthio, alkylsulfinyl, and alkylsulfonyl. Exemplary aryl groups include phenyl, naphthyl, tetrahydronaphthyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-methylphenyl, 4-methoxyphenyl, 3-trifluoromethylphenyl, 4-nitrophenyl, and the like.

The term "heteroaryl" is defined herein as a monocyclic or bicyclic ring system containing one or two aromatic rings and containing at least one nitrogen, oxygen, or sulfur atom in an aromatic ring, and which can be unsubstituted or substituted, for example, with one or more, and in particular one to three, substituents, like halo, alkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkyl, nitro, amino, alkylamino, acylamino, alkylthio, alkylsulfinyl, and alkylsulfonyl. Examples of heteroaryl groups include thienyl, furyl, pyridyl, oxazolyl, quinolyl, isoquinolyl, indolyl, triazolyl, isothiazolyl, isoxazolyl, imidazolyl, benzothiazolyl, pyrazinyl, pyrimidinyl, thiazolyl, and thiadiazolyl.

The term "alkoxyalkyl" is defined as an alkyl group wherein a hydrogen has been replaced by an alkoxy group. The term "(alkylthio)alkyl" is defined similarly as alkoxyalkyl, except a sulfur atom, rather than an oxygen atom, is present.

The term "hydroxy" is defined as -OH.

- 7 -

The term "alkoxy" is defined as -OR, wherein R is alkyl.

The term "hydroxyalkyl" is defined as a hydroxy group appended to an alkyl group.

5           The term "amino" is defined as -NH<sub>2</sub>, and the term "alkylamino" is defined as -NR<sub>2</sub>, wherein at least one R is alkyl and the second R is alkyl or hydrogen.

10           The term "acylamino" is defined as RC(=O)N, wherein R is alkyl or aryl.

The term "alkylthio" is defined as -SR, wherein R is alkyl.

The term "alkylsulfinyl" is defined as R-SO<sub>2</sub>, wherein R is alkyl.

15           The term "alkylsulfonyl" is defined as R-SO<sub>3</sub>, wherein R is alkyl.

The term "nitro" is defined as -NO<sub>2</sub>.

The term "trifluoromethyl" is defined as -CF<sub>3</sub>.

20           The term "trifluoromethoxy" is defined as -OCF<sub>3</sub>.


The term "cyano" is defined as -CN.

25           In a preferred embodiment, R<sup>1</sup> is selected from the group consisting of hydrogen, C<sub>1-6</sub>alkyl, haloC<sub>1-6</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkylC<sub>1-3</sub>alkyl, arylC<sub>2-3</sub>alkyl, and heteroarylC<sub>2-3</sub>alkyl.

In a preferred embodiment, R<sup>2</sup> is the optionally substituted bicyclic ring system




- 8 -



5

wherein the bicyclic ring can represent,  
for example, naphthalene or indene, or a hetero-  
cycle, such as benzoxazole, benzothiazole, benzisox-  
azole, benzimidazole, quinoline, indole, benzothio-  
phene, or benzofuran, or

10




15

wherein  $n$  is an integer 1 or 2, and  $X$ , independ-  
ently, are  $C(R^a)_2$ ,  $O$ ,  $S$ , or  $NR^a$ . The bicyclic ring  
comprising the  $R^2$  substituent typically is attached  
to the rest of the molecule by a phenyl ring carbon  
atom.

20

In a preferred group of compounds of for-  
mula (I),  $R^2$  is represented by an optionally sub-  
stituted bicyclic ring

25




30


- 9 -

wherein n is 1 or 2, and X, independently, are CH<sub>2</sub> or O. Especially preferred R<sup>2</sup> substituents include

5




10



, and

15



20

Within this particular group of compounds, nonlimiting examples of substituents for the bicyclic ring include halogen (e.g., chlorine), C<sub>1-3</sub>alkyl (e.g., methyl, ethyl, or i-propyl), OR<sup>a</sup> (e.g., methoxy, ethoxy, or hydroxy), CO<sub>2</sub>R<sup>a</sup>, halomethyl or halomethoxy (e.g., trifluoromethyl or trifluoromethoxy), cyano, 25 nitro, and NR<sup>a</sup>R<sup>b</sup>.


30

In another preferred embodiment, R<sup>4</sup> is selected from the group consisting of aryl, Het, OR<sup>a</sup>, C(=O)OR<sup>a</sup>, C<sub>1-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>, OC(=O)R<sup>a</sup>, C(=O)R<sup>a</sup>, NR<sup>a</sup>R<sup>b</sup>, C<sub>3-8</sub>cycloalkyl, C<sub>3-8</sub>cycloalkyl, C(=O)NR<sup>a</sup>R<sup>b</sup>, CF<sub>3</sub>, OCF<sub>3</sub>, CN, SO<sub>2</sub>NR<sup>a</sup>R<sup>b</sup>, OC<sub>2-4</sub>alkyleneNR<sup>a</sup>R<sup>b</sup>, and C(=O)NR<sup>a</sup>R<sup>c</sup>, or R<sup>4</sup> and R<sup>5</sup> are taken together with the carbon atoms to which they are attached to form a 5- or 6-membered

- 10 -

ring, saturated or partially or fully saturated, optionally substituted and optionally containing one or two heteroatoms.

An especially preferred subclass of compounds within the general scope of formula (I) is represented by compounds of formula (II)



(II)

and pharmaceutically acceptable salts and solvates (e.g., hydrates) thereof.

Compounds of formula (I) can contain one or more asymmetric center, and, therefore, can exist as stereoisomers. The present invention includes both mixtures and separate individual stereoisomers of the compounds of formula (I). Compounds of formula (I) also can exist in tautomeric forms, and the invention includes both mixtures and separate individual tautomers thereof.

Pharmaceutically acceptable salts of the compounds of formula (I) can be acid addition salts formed with pharmaceutically acceptable acids. Examples of suitable salts include, but are not limited to, the hydrochloride, hydrobromide, sul-

- 11 -

fate, bisulfate, phosphate, hydrogen phosphate, acetate, benzoate, succinate, fumarate, maleate, lactate, citrate, tartrate, gluconate, methanesulfonate, benzenesulfonate, and p-toluenesulfonate salts. The compounds of the formula (I) also can provide pharmaceutically acceptable metal salts, in particular alkali metal salts and alkaline earth metal salts, with bases. Examples include the sodium, potassium, magnesium, and calcium salts.

Compounds of the present invention are potent and selective inhibitors of cGMP-specific PDE5. Thus, compounds of formula (I) are of interest for use in therapy, specifically for the treatment of a variety of conditions where selective inhibition of PDE5 is considered to be beneficial.

Phosphodiesterases (PDEs) catalyze the hydrolysis of cyclic nucleotides, such as cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP). The PDEs have been classified into at least seven isoenzyme families and are present in many tissues (J.A. Beavo, *Physiol. Rev.*, 75, p. 725 (1995)).

PDE5 inhibition is a particularly attractive target. A potent and selective inhibitor of PDE5 provides vasodilating, relaxing, and diuretic effects, all of which are beneficial in the treatment of various disease states. Research in this area has led to several classes of inhibitors based on the cGMP basic structure (E. Sybertz et al., *Expert. Opin. Ther. Pat.*, 7, p. 631 (1997)).

The biochemical, physiological, and clinical effects of PDE5 inhibitors therefore suggest their utility in a variety of disease states in

- 12 -

which modulation of smooth muscle, renal, hemostatic, inflammatory, and/or endocrine function is desirable. The compounds of formula (I), therefore, have utility in the treatment of a number of disorders, including stable, unstable, and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, acute respiratory distress syndrome, acute and chronic renal failure, atherosclerosis, conditions of reduced blood vessel patency (e.g., postpercutaneous transluminal coronary or carotid angioplasty, or post-bypass surgery graft stenosis), peripheral vascular disease, vascular disorders, such as Raynaud's disease, thrombocytopenia, inflammatory diseases, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, osteoporosis, preterm labor, benign prostatic hypertrophy, peptic ulcer, male erectile dysfunction, female sexual dysfunction, and diseases characterized by disorders of gut motility (e.g., irritable bowel syndrome).

An especially important use is the treatment of male erectile dysfunction, which is one form of impotence and is a common medical problem. Impotence can be defined as a lack of power, in the male, to copulate, and can involve an inability to achieve penile erection or ejaculation, or both. The incidence of erectile dysfunction increases with age, with about 50% of men over the age of 40 suffering from some degree of erectile dysfunction.

In addition, a further important use is the treatment of female arousal disorder, also termed female sexual arousal disorder. Female arousal disorders are defined as a recurrent inability

- 13 -

ity to attain or maintain an adequate lubrication/-  
swelling response of sexual excitement until comple-  
tion of sexual activity. The arousal response con-  
sists of vasocongestion in the pelvis, vaginal lu-  
5 brication, and expansion and swelling of external  
genitalia.

It is envisioned, therefore, that com-  
pounds of formula (I) are useful in the treatment of  
male erectile dysfunction and female sexual arousal  
10 disorder. Thus, the present invention concerns the  
use of compounds of formula (I), or a pharmaceuti-  
cally acceptable salt thereof, or a pharmaceutical  
composition containing either entity, for the manu-  
facture of a medicament for the curative or prophy-  
15 lactic treatment of erectile dysfunction in a male  
animal and sexual arousal disorder in a female ani-  
mal, including humans.

The term "treatment" includes preventing,  
lowering, stopping, or reversing the progression or  
20 severity of the condition or symptoms being treated.  
As such, the term "treatment" includes both medical  
therapeutic and/or prophylactic administration, as  
appropriate.

It also is understood that "a compound of  
25 formula (I)," or a physiologically acceptable salt  
or solvate thereof, can be administered as the neat  
compound, or as a pharmaceutical composition con-  
taining either entity.

Although the compounds of the invention  
30 are envisioned primarily for the treatment of sexual  
dysfunction in humans, such as male erectile dys-  
function and female sexual arousal disorder, they  
also can be used for the treatment of other disease

- 14 -

states. A further aspect of the present invention, therefore, is providing a compound of formula (I) for use in the treatment of stable, unstable, and variant (Prinzmetal) angina, hypertension, pulmonary  
5 hypertension, chronic obstructive pulmonary disease, congestive heart failure, acute respiratory distress syndrome, acute and chronic renal failure, atherosclerosis, conditions of reduced blood vessel patency (e.g., post-PTCA or post-bypass graft stenosis),  
10 peripheral vascular disease, vascular disorders such as Raynaud's disease, thrombocythemia, inflammatory diseases, prophylaxis of myocardial infarction, prophylaxis of stroke, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma,  
15 osteoporosis, preterm labor, benign prostatic hypertrophy, male and female erectile dysfunction, or diseases characterized by disorders of gut motility (e.g., IBS).

According to another aspect of the present  
20 invention, there is provided the use of a compound of formula (I) for the manufacture of a medicament for the treatment of the above-noted conditions and disorders.

In a further aspect, the present invention  
25 provides a method of treating the above-noted conditions and disorders in a human or nonhuman animal body which comprises administering to said body a therapeutically effective amount of a compound of formula (I).

30 Compounds of the invention can be administered by any suitable route, for example by oral, buccal, inhalation, sublingual, rectal, vaginal, transurethral, nasal, topical, percutaneous, i.e.,

- 15 -

transdermal, or parenteral (including intravenous, intramuscular, subcutaneous, and intracoronary) administration. Parenteral administration can be accomplished using a needle and syringe, or using a  
5 high pressure technique, like POWDERJECT™.

Oral administration of a compound of the invention is the preferred route. Oral administration is the most convenient and avoids the disadvantages associated with other routes of administration. For patients suffering from a swallowing disorder or from impairment of drug absorption after oral administration, the drug can be administered parenterally, e.g., sublingually or buccally.  
10

Compounds and pharmaceutical compositions suitable for use in the present invention include those wherein the active ingredient is administered in an effective amount to achieve its intended purpose. More specifically, a "therapeutically effective amount" means an amount effective to prevent development of, or to alleviate the existing symptoms of, the subject being treated. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.  
15  
20

A "therapeutically effective dose" refers to that amount of the compound that results in achieving the desired effect. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population). The dose ratio between  
25  
30



- 16 -

toxic and therapeutic effects is the therapeutic index, which is expressed as the ratio between LD<sub>50</sub> and ED<sub>50</sub>. Compounds which exhibit high therapeutic indices are preferred. The data obtained from such data can be used in formulating a range of dosage for use in humans. The dosage of such compounds preferably lies within a range of circulating concentrations that include the ED<sub>50</sub> with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed, and the route of administration utilized.

The exact formulation, route of administration, and dosage can be chosen by the individual physician in view of the patient's condition. Dosage amount and intervals can be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain therapeutic effects.

The amount of composition administered is dependent on the subject being treated, the subject's weight, the severity of the affliction, the manner of administration, and the judgment of the prescribing physician.

Specifically, for administration to a human in the curative or prophylactic treatment of the conditions and disorders identified above, oral dosages of a compound of formula (I) generally are about 0.5 to about 1000 mg daily for an average adult patient (70 kg). Thus, for a typical adult patient, individual tablets or capsules contain 0.2 to 500 mg of active compound, in a suitable pharmaceutically acceptable vehicle or carrier, for administration in single or multiple doses, once or sev-

- 17 -

eral times per day. Dosages for intravenous, buccal, or sublingual administration typically are 0.1 to 500 mg per single dose as required. In practice, the physician determines the actual dosing regimen which is most suitable for an individual patient, and the dosage varies with the age, weight, and response of the particular patient. The above dosages are exemplary of the average case, but there can be individual instances wherein higher or lower dosages are merited, and such are within the scope of this invention.

For human use, a compound of the formula (I) can be administered alone, but generally is administered in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. Pharmaceutical compositions for use in accordance with the present invention thus can be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries that facilitate processing of compounds of formula (I) into preparations which can be used pharmaceutically.

These pharmaceutical compositions can be manufactured in a conventional manner, e.g., by conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen. When a therapeutically effective amount of a compound of the present invention is administered orally, the composition typically is in the form of a tablet, capsule, powder, solution, or

- 18 -

elixir. When administered in tablet form, the composition can additionally contain a solid carrier, such as a gelatin or an adjuvant. The tablet, capsule, and powder contain about 5% to about 95% compound of the present invention, and preferably from about 25% to about 90% compound of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, or oils of animal or plant origin can be added. The liquid form of the composition can further contain physiological saline solution, dextrose or other saccharide solutions, or glycols. When administered in liquid form, the composition contains about 0.5% to about 90% by weight of a compound of the present invention, and preferably about 1% to about 50% of a compound of the present invention.

When a therapeutically effective amount of a compound of the present invention is administered by intravenous, cutaneous, or subcutaneous injection, the composition is in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred composition for intravenous, cutaneous, or subcutaneous injection typically contains, in addition to a compound of the present invention, an isotonic vehicle.

For oral administration, the compounds readily can be formulated by combining a compound of formula (I) with pharmaceutically acceptable carriers well known in the art. Such carriers enable the present compounds to be formulated as tablets,

- 19 -

pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by adding  
5 a compound of formula (I) with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, for  
10 example, fillers and cellulose preparations. If desired, disintegrating agents can be added.

For administration by inhalation, compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the  
15 use of a suitable propellant. In the case of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin, for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a  
20 suitable powder base such as lactose or starch.

The compounds can be formulated for parenteral administration by injection, e.g., by  
25 bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, e.g., in ampules or in multidose containers, with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and can contain  
30 formulatory agents such as suspending, stabilizing, and/or dispersing agents.

- 20 -

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils or synthetic fatty acid esters. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension. Optionally, the suspension also can contain suitable stabilizers or agents that increase the solubility of the compounds and allow for the preparation of highly concentrated solutions. Alternatively, a present composition can be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

Compounds of the present invention also can be formulated in rectal compositions, such as suppositories or retention enemas, e.g., containing conventional suppository bases. In addition to the formulations described previously, the compounds also can be formulated as a depot preparation. Such long-acting formulations can be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Many of the compounds of the present invention can be provided as salts with pharmaceutically compatible counterions. Such pharmaceutically

- 21 -

acceptable base addition salts are those salts that retain the biological effectiveness and properties of the free acids, and that are obtained by reaction with suitable inorganic or organic bases.

5                   In particular, a compound of formula (I) can be administered orally, buccally, or sublingually in the form of tablets containing excipients, such as starch or lactose, or in capsules or ovules, either alone or in admixture with excipients, or in  
10                   the form of elixirs or suspensions containing flavoring or coloring agents. Such liquid preparations can be prepared with pharmaceutically acceptable additives, such as suspending agents. A compound also can be injected parenterally, for example,  
15                   intravenously, intramuscularly, subcutaneously, or intracoronarily. For parenteral administration, the compound is best used in the form of a sterile aqueous solution which can contain other substances, for example, salts, or monosaccharides, such as mannitol  
20                   or glucose, to make the solution isotonic with blood.

                  For veterinary use, a compound of formula (I) or a nontoxic salt thereof, is administered as a suitably acceptable formulation in accordance with  
25                   normal veterinary practice. The veterinarian can readily determine the dosing regimen and route of administration that is most appropriate for a particular animal.


                  Thus, the invention provides in a further  
30                   aspect a pharmaceutical composition comprising a compound of the formula (I), together with a pharmaceutically acceptable diluent or carrier therefor. There is further provided by the present invention a

- 22 -

process of preparing a pharmaceutical composition comprising a compound of formula (I), which process comprises mixing a compound of formula (I), together with a pharmaceutically acceptable diluent or carrier therefor.

In a particular embodiment, the invention includes a pharmaceutical composition for the curative or prophylactic treatment of erectile dysfunction in a male animal, or arousal disorder in a female animal, including humans, comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable diluent or carrier.

Compounds of formula (I) can be prepared by any suitable method known in the art, or by the following processes which form part of the present invention. In the methods below,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ , and  $R^5$  are as defined in structural formula (I) above. In particular, Daugan U.S. Patent No. 5,859,006, incorporated herein by reference, discloses preparation of a compound of structural formula (III).




(III)

Daugan U.S. Patent No. 5,859,006 teaches the preparation of the compound of structural formula (III),

- 23 -


wherein R<sup>0</sup> is H, beginning with a tryptophan ester having the structural formula (IV):



(IV)

D-Tryptophan ester

The compounds of structural formula (I) can be prepared in an analogous manner as a compound of structural formula (III) using an appropriately substituted tryptophan ester of structural formula (V):



(V)

Many substituted tryptophan esters of structural formula (V) are commercially available, and, if necessary, can be converted to other substituted esters. Substituted tryptophan esters also can be prepared, for example, from substituted indoles, as set forth in S. Wagaw et al., *J. Amer. Chem. Soc.*, 21, p. 10251 (1999) and M.P. Moyer et al., *J. Org. Chem.*, 51, p. 5106 (1986). These substituted tryptophan esters can be used in the syn-



- 24 -

thetic methods disclosed in Daugan U.S. Patent No. 5,859,006 to provide compositions of structural formula (I).

It should be understood that protecting groups can be utilized in accordance with general principles of organic synthetic chemistry to provide compounds of structural formula (I). Protecting compounds and protecting groups, like benzyl chloroformate and trichloroethyl chloroformate, are well known to persons skilled in the art, for example, see T.W. Greene et al., "Protective Groups in Organic Synthesis, Third Edition," John Wiley and Sons, Inc., NY, NY (1999). These protecting groups are removed when necessary by appropriate basic, acidic, or hydrogenolytic conditions known to persons skilled in the art. Accordingly, compounds of structural formula (I) not specifically exemplified herein can be prepared by persons skilled in the art.

In addition, compounds of formula (I) can be converted to other compounds of formula (I). Thus, for example, a particular  $R^4$  substituent can be interconverted to prepare another suitably substituted compound of formula (I). Examples of appropriate interconversions include, but are not limited to, nitro to amino,  $OR^a$  to hydroxy by suitable means (e.g., using a reducing agent such as  $SnCl_2$  or a palladium catalyst, such as palladium-on-carbon), or amino to substituted amino, such as acylamino or sulphonylamino, using standard acylating or sulfonylating conditions. In cases wherein  $R^1$  represents a substituted bicyclic system, suitable interconversion can involve removal of a substituent, such as

- 25 -

by treatment with a palladium catalyst whereby, for example, a benzyl substituent is removed from a suitable bicyclic system.

Compounds of formula (I) can be prepared by the method above as individual stereoisomers from the appropriate stereoisomer of formula (IV) or as a racemic mixture from the appropriate racemic compound of formula (IV). Individual stereoisomers of the compounds of the invention can be prepared from racemates by resolution using methods known in the art for the separation of racemic mixtures into their constituent stereoisomers, for example, using HPLC on a chiral column, such as Hypersil naphthyl urea, or using separation of salts of stereoisomers. Compounds of the invention can be isolated in association with solvent molecules by crystallization from, or evaporation of, an appropriate solvent.


The pharmaceutically acceptable acid addition salts of the compounds of formula (I) that contain a basic center can be prepared in a conventional manner. For example, a solution of the free base can be treated with a suitable acid, either neat or in a suitable solution, and the resulting salt isolated either by filtration or by evaporation under vacuum of the reaction solvent. Pharmaceutically acceptable base addition salts can be obtained in an analogous manner by treating a solution of a compound of formula (I) with a suitable base. Both types of salt can be formed or interconverted using ion-exchange resin techniques. Thus, according to a further aspect of the invention, a method for preparing a compound of formula (I) or a salt or solvate (e.g., hydrate) is provided, followed by (i)

- 26 -

salt formation, or (ii) solvate (e.g., hydrate) formation.

The following abbreviations are used hereafter in the accompanying examples: rt (room temperature), min (minute), h (hour), g (gram), mmol (millimole), m.p. (melting point), eq (equivalents), aq (aqueous), L (liter), mL (milliliter),  $\mu$ L (microliter), DMSO (dimethyl sulfoxide),  $\text{CH}_2\text{Cl}_2$  (dichloromethane), IPA (isopropyl alcohol), TFA (trifluoroacetic acid), NaOH (sodium hydroxide), TsOH (p-toluenesulfonic acid), Me (methyl), Et (ethyl), EtOH (ethanol), MeOH (methanol), DMF (dimethylformamide),  $\text{Et}_3\text{N}$  (triethylamine),  $\text{MeNH}_2$  (methylamine), HOAc (acetic acid),  $\text{Ac}_2\text{O}$  (acetic anhydride), Ac ( $\text{C}(=\text{O})\text{CH}_3$ ), and THF (tetrahydrofuran).

#### Preparation of Example 1





Example 1


Example 1 was prepared from 5-hydroxy-DL-tryptophan as depicted in the following synthetic scheme. 5-Hydroxy-DL-tryptophan is a commercially

- 27 -


available compound from Aldrich Chemical Co., Milwaukee, WI.



Intermediate 1




0°C to rt  
12% *cis*  
12% *trans*




Intermediate 2

- 28 -



Intermediate 3



64% for two steps

Example 1

Intermediate 1Preparation of 5-hydroxy-DL-tryptophan  
methyl ester hydrochloride

30

Thionyl chloride (2.14 mL, 29.4 mmol) was added dropwise to a slurry of 5-hydroxy-DL-tryptophan (3.8 g, 17.3 mmol) in anhydrous methanol (40 mL) at 0°C under a nitrogen blanket. The resulting

- 29 -

mixture was warmed slowly to room temperature and stirred for 17 hours. Then the solvent was removed under reduced pressure to provide 5-hydroxy-DL-tryptophan methyl ester hydrochloride (Intermediate 1) as a light brown foam (5.0 g, 100%):  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  10.78 (bs, 1H), 8.62-8.25 (m, 3H), 7.18-7.11 (m, 2H), 6.79 (s, 1H), 6.67-6.61 (m, 1H), 4.18 (m, 1H), 3.69 (s, 3H), 3.25-3.12 (m, 2H).

## Intermediate 2

### Preparation of a *cis*- $\beta$ -Carboline

A mixture of Intermediate 1 (2.7 g, 10 mmol) and piperonal (1.8 g, 12 mmol) in IPA (27 mL) was heated at reflux under a nitrogen blanket for 8 hours, during which time a precipitate formed (after about 4 hours). The resulting slurry was cooled to room temperature, vacuum filtered, and the solid was washed with IPA (2 x 5 mL). The filtrate solvent was removed under reduced pressure, then the residue was diluted with ethyl acetate (60 mL) and neutralized with saturated sodium bicarbonate ( $\text{NaHCO}_3$ ) solution (20 mL). The basic aqueous layer was extracted with ethyl acetate, then the combined organic extracts were concentrated under reduced pressure to provide a yellow solid residue. The residue was purified by column chromatography, eluting with methylene chloride/ethyl acetate (4:1), to provide Intermediate 2 as a yellow solid (420 mg, 12%): TLC  $R_f$  (4:1 methylene chloride/ethyl acetate)=0.55;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.31 (bs, 1H), 7.07 (d,  $J=8.5$  Hz, 1H), 6.92-6.68 (m, 5H), 5.95 (s, 2H), 5.14 (s, 1H), 3.97-3.91 (m, 1H), 3.81 (s, 3H), 3.19-3.08

- 30 -

(m, 1H), 3.00-2.88 (m 1H). The *trans* carboline also was obtained as a yellow solid, but was not characterized (420 mg, 12%): TLC  $R_f$  (4:1 methylene chloride/ethyl acetate)=0.20.

5

### Intermediate 3

#### Preparation of (+/-)-*cis*-2-chloroacetyl- $\beta$ -carboline

10

Chloroacetyl chloride (0.36 mL, 4.52 mmol) was added dropwise to a mixture of Intermediate 2 (500 mg, 1.28 mmol) and triethylamine (0.63 mL, 4.54 mmol) in methylene chloride (20 mL) at 0°C under a nitrogen blanket, and the resulting mixture was stirred at 5°C for 2.5 hours. The resulting brown solution was diluted with methylene chloride (40 mL) and washed successively with saturated NaHCO<sub>3</sub> (15 mL) and brine (10 mL). The organic layer was dried over sodium sulfate (Na<sub>2</sub>SO<sub>4</sub>), then the solvent was removed under reduced pressure to provide Intermediate 3 as a thick yellow oil (0.95 g), which was used without purification: TLC  $R_f$  (4:1 methylene chloride/ethyl acetate)=0.92.

15

20

25

### Example 1

#### Preparation of (+/-, *cis*)-6-Benzo[1,3]dioxol-5-yl-10-hydroxy-2-methyl-2,3,6,7,12,12a-hexahydro-pyrazino[1',2':1,6)pyrido[3,4-b]indole-1,4-dione

30

A mixture of crude Intermediate 3 (0.95 g), methylamine (4.5 mL, 2.0M in THF, 9 mmol), and methanol (8 mL) was heated at 50°C under a nitrogen blanket for 5 hours, after which the resulting mix-


35

- 31 -


ture was heated for 17 hours at 40°C. The resulting orange slurry was cooled to room temperature, then the precipitate was collected by vacuum filtration. The solid was washed with methanol (5 x 2 mL), slurried in diethyl ether for 5 hours, then the solid was collected by vacuum filtration. The solid next was dried in a vacuum oven at 70°C for 24 hours to provide the compound of Example 1 as a white powder (0.33 g, 64% for two steps). The desired *cis* isomer was confirmed by an NOE difference experiment (1.0% enhancement): mp 322-328°C; TLC R<sub>f</sub> (4:1:0.4 methylene chloride/ethyl acetate/methanol)=0.55; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 10.70 (s, 1H), 8.67 (s, 1H), 7.08 (d, J=8.6 Hz, 1H), 6.85-6.75 (m, 3H), 6.60-6.55 (m, 1H), 6.09 (s, 1H), 5.93 (s, 2H), 4.42-4.33 (m, 1H), 4.18 (d, J=17.0 Hz, 1H), 3.93 (d, J=17.0 Hz, 1H), 3.45-3.32 (m, 1H), 3.00-2.88 (m, 4H); <sup>13</sup>C NMR (75 MHz, DMSO-d<sub>6</sub>): δ 166.8, 166.6, 150.6, 146.9, 145.9, 137.0, 134.2, 130.6, 136.4, 119.1, 111.6, 111.2, 107.9, 106.8, 103.9, 102.1, 100.8, 55.4, 55.1, 51.4, 32.8, 23.0 ppm; CI MS (methane) m/z 406 [M+H]<sup>+</sup>. Anal. Calcd. for C<sub>22</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub>; C, 65.18; H, 4.72; N, 10.37. Found: C, 64.47; H, 5.04; N, 10.24.



- 32 -


Preparation of Example 2

Example 2 was prepared from 5-methoxy-DL-tryptophan as depicted in the following synthetic scheme. 5-Methoxy-DL-tryptophan is a commercially available compound from Aldrich Chemical Co., Milwaukee, WI.




Intermediate 4

- 33 -




Intermediate 5



Intermediate 6

- 34 -



Example 2

#### Intermediate 4

#### Preparation of 5-Methoxy-DL-tryptophan methyl ester

Thionyl chloride (1.0 mL, 13.7 mmol) was added dropwise to a slurry of 5-methoxy-DL-tryptophan (2.0 g, 8.54 mmol) in anhydrous methanol (22 mL) at 0°C under a nitrogen blanket. The resulting solution was warmed slowly to room temperature, stirred for 17 hours, then the solvent was removed under reduced pressure. The resulting residue was dissolved in water (20 mL), neutralized with saturated NaHCO<sub>3</sub> (10 mL), and extracted twice with methylene chloride. The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and the solvent was removed under reduced pressure to provide 5-methoxy-DL-tryptophan methyl ester (Intermediate 4) as a light brown solid (1.55 g, 73%): <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.93 (bs, 1H), 7.30-7.25 (m, 1H), 7.09-7.03 (m, 1H), 6.90-6.83 (m, 1H), 3.90-3.78 (m, 4H), 3.72 (s, 3H), 3.30-3.20 (m, 1H), 3.10-2.98 (m, 1H).

- 35 -

Intermediate 5Preparation of the (+/-)-cis- $\beta$ -Carboline

Trifluoroacetic acid (0.65 mL, 8.48 mmol) was added to a mixture of Intermediate 4 (1.6 g, 6.45 mmol) and piperonal (1.17 g, 7.74 mmol) in methylene chloride (30 mL) at 0°C under a nitrogen blanket. The resulting mixture was warmed to room temperature then stirred for 48 hours. The reaction mixture was diluted with methylene chloride (70 mL), then neutralized with saturated NaHCO<sub>3</sub> (15 mL). The organic layer was concentrated under reduced pressure, and the residue was purified by flash column chromatography, eluting with methylene chloride/ethyl acetate (8:1), to provide Intermediate 5 as a yellow solid (882 mg, 36%): TLC R<sub>f</sub> (4:1 methylene chloride/ethyl acetate)=0.69; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  7.32 (bs, 1H), 7.14 (d, J=8.7 Hz, 1H), 7.00 (s, 1H), 6.93-6.80 (m, 4H), 5.97 (s, 2H), 5.18 (s, 1H), 4.02-3.94 (m, 1H), 3.88 (s, 3H), 3.84 (s, 3H), 3.27-3.16 (m, 1H), 3.07-2.96 (m, 1H). The *trans* carboline also was obtained as a yellow solid, but was not characterized (850 mg, 35%): TLC R<sub>f</sub> (4:1: methylene chloride/ethyl acetate)=0.40.

Intermediate 6Preparation of the (+/-)-cis-2-Chloroacetyl- $\beta$ -carboline

Chloroacetyl chloride (0.21 mL, 2.60 mmol) was added dropwise to a mixture of Intermediate 5 (880 mg, 2.17 mmol) and triethylamine (0.36 mL, 2.60 mmol) in methylene chloride (18 mL) at 0°C under a

- 36 -

nitrogen blanket, and the resulting mixture was stirred at 0°C for 2 hours. The resulting solution then was diluted with methylene chloride (60 mL), washed with saturated NaHCO<sub>3</sub> (10 mL), dried over sodium sulfate, filtered, and the solvent was removed under reduced pressure to provide Intermediate 6 as a yellow foam which was used without purification (1.14 g): TLC R<sub>f</sub> (4:1 methylene chloride/ethyl acetate)=0.92.

### Example 2


Preparation of (trans)-6-(1,3-Benzodioxol-5-yl)-10-methoxy-2-methyl-2,3,6,7,12,12a-hexahydro-pyrazino[1',2':1,6]pyrido[3,4-b]indole-1,4-dione

A mixture of crude Intermediate 6 (1.14 g, 2.17 mmol) and methylamine (5 mL, 2.0 M in THF, 10 mmol) in methanol (12 mL) was heated at 50°C under a nitrogen blanket for 6 hours, after which the mixture was stirred at room temperature for 17 hours. The resulting precipitate was isolated by vacuum filtration, then the solid was washed with methanol (2 x 5 mL) and dried in a vacuum oven at 60°C for 24 hours to provide Example 2 as a white powder (0.65 g, 71% over two steps): mp 259-261°C; TLC R<sub>f</sub> (4:1 methylene chloride/ethyl acetate)=0.35; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 10.81 (s, 1H), 7.18 (d, J=8.8 Hz, 1H), 7.09 (d, J=1.5 Hz, 1H), 6.85 (s, 1H), 6.78 (s, 2H), 6.72-6.62 (m, 1H), 6.10 (s, 1H), 5.93 (s, 2H), 4.43-4.32 (m, 1H), 4.18 (d, J=17.2 Hz, 1H), 3.94 (d, J=17.2 Hz, 1H), 3.57-3.45 (m 1H), 3.01-2.82 (m, 4H): <sup>13</sup>C NMR (75 MHz, DMSO-d<sub>6</sub>) δ: 166.8, 166.6, 153.4, 147.0, 146.0, 137.0, 134.5, 131.1, 126.1, 119.2,

- 37 -



111.9, 111.2, 107.9, 106.9, 104.6, 100.8, 100.2, 55.5, 55.3, 55.2, 51.4, 32.8, 23.2 ppm; CI MS (methane)  $m/z$  420  $[C_{23}H_{21}N_3O_5+H]^+$ . Anal. Calcd. for  $C_{23}H_{21}N_3O_5$ : C, 65.86; H, 5.05; N, 10.02. Found: C, 65.75; H, 4.95; N, 9.91. The relative stereochemistry of Example 2 was confirmed to be *cis* isomer by an NOE difference experiment (DMSO- $d_6$ ): a positive NOE enhancement (2.3%) from the C12a proton at 4.37 ppm to the C6 proton at 6.12 ppm.

### Preparation of Example 3



Example 3 was prepared from 6-methoxy-D-tryptophan ethyl ester as depicted in the following synthetic scheme. 6-Methoxy-D-tryptophan ethyl ester is a commercially available compound from Aldrich Chemical Co., Milwaukee, WI.

- 38 -





Intermediate 7

25

- 39 -



5



10

Intermediate 8

15



20

Example 3

Intermediate 7Preparation of a *cis*- $\beta$ -Carboline

25

30

Trifluoroacetic acid (0.295 mL, 3.82 mmol) was added to a mixture of 6-methoxy-D-tryptophan ethyl ester (500 mg, 1.91 mmol) and piperonal (573 mg, 3.82 mmol) in methylene chloride (15 mL) at 0°C under a nitrogen blanket, after which the mixture was warmed to room temperature overnight, then stirred for 3 days. The reaction mixture then was diluted with methylene chloride (30 mL) and neutral-



- 40 -

ized with saturated  $\text{NaHCO}_3$  (5 mL). The organic layer then was washed with brine (5 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and the solvent was removed under reduced pressure to provide a brown solid residue. The residue was purified by flash column chromatography, eluting with methylene chloride/ethyl acetate (10:1), to provide Intermediate 7 as a yellow solid (120 mg, 15%): TLC  $R_f$  (10:1 methylene chloride/ethyl acetate)=0.57;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.42-7.30 (m, 2H), 6.90-6.70 (m, 6H), 5.94 (s, 2H), 5.13 (s, 1H), 4.31-4.19 (m, 2H), 3.93-3.86 (m, 1H), 3.79 (s, 3H), 3.20-3.10 (m, 1H), 3.00-2.89 (m, 1H), 1.35 (t,  $J=8.5$  Hz, 3H). The *trans*- $\beta$ -carboline also was obtained as a yellow solid, but was not characterized (160 mg, 20%): TLC  $R_f$  (10:1 methylene chloride/ethyl acetate)=0.29.

### Intermediate 8

#### Preparation of a *cis*-2-Chloroacetyl- $\beta$ -carboline

Chloroacetyl chloride (0.030 mL, 0.373 mmol) was added dropwise to a mixture of Intermediate 7 (120 mg, 0.287 mmol) and triethylamine (0.052 mL, 0.373 mmol) in methylene chloride (5 mL) at  $0^\circ\text{C}$  under a nitrogen blanket, and the resulting mixture was stirred at  $0^\circ\text{C}$  for 1.5 hours. The brown solution then was diluted with methylene chloride (30 mL), washed with saturated  $\text{NaHCO}_3$  (5 mL) and brine (5 mL), and the solvent was removed under reduced pressure to provide Intermediate 8 as a yellow solid which was used without further purification (179 mg): TLC  $R_f$  (4:1 methylene chloride/ethyl acetate)=0.92.

- 41 -

Example 3

Preparation of (6R,12aR)-6-Benzo[1,3]dioxol-5-yl-9-methoxy-2-methyl-2,3,6,7,12,12a-hexahydropyrazino-[1',2':1,6]pyrido[3,4-b]indole-1,4-dione

5


A mixture of crude Intermediate 8 (179 mg) and methylamine (0.5 mL, 2.0M in THF, 1.0 mmol) in THF (3 mL) and methanol (5 mL) was heated at 50°C under a nitrogen blanket for 6 hours. A second portion of methylamine (0.25 mL, 0.5 mmol) was added, and the resulting mixture was stirred at 40°C for an additional 16 hours. The resulting orange slurry was cooled to room temperature, then the mixture was concentrated to about 5 mL. The resulting precipitate was isolated by vacuum filtration. The solid then was washed with methanol (3 x 2 mL) and dried in a vacuum oven at 70°C to provide Example 3 as an off-white powder (0.085 g, 71% over two steps): mp 288-293°C; TLC R<sub>f</sub> (4:1 methylene chloride/ethyl acetate)=0.38; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 10.81 (s, 1H), 7.42 (d, J=8.6 Hz, 1H), 6.84 (s, 1H), 6.80 (s, 1H), 6.77 (s, 2H), 6.65 (dd, J=8.6, 2.1 Hz, 1H), 6.09 (s, 1H), 5.92 (s, 2H), 4.38 (dd, J=11.6, 4.3 Hz, 1H), 4.17 (d, J=19.4 Hz, 1H), 3.93 (d, J=19.4 Hz, 1H), 3.73 (s, 3H), 3.46 (dd, J=15.9, 4.5 Hz, 1H), 3.00-2.87 (m, 4H); API m/z 420 [C<sub>23</sub>H<sub>21</sub>N<sub>3</sub>O<sub>5</sub>+H]<sup>+</sup>; [α]<sub>D</sub><sup>25°C</sup>=+136.7° (c=0.25, DMSO). Anal. Calcd. for C<sub>23</sub>H<sub>21</sub>N<sub>3</sub>O<sub>5</sub>; C, 65.86; H, 5.05; N, 10.02. Found: C, 65.96; H, 4.97; N, 9.91. The stereochemistry of Example 3 was confirmed to be the desired *cis* isomer by a series of NOE difference experiments: a positive NOE enhancement from the C12a proton at 4.38 ppm to the C6 proton at 6.09 ppm; a

30


- 42 -

positive NOE enhancement from the C6 proton at 6.09 ppm to the C12a proton at 4.38 ppm. Chiral HPLC analysis (Chiralcel OD Column, 250 x 4.6 mm, Retention Time=20.9 minutes; 1:1 isopropanol/hexanes; flow=1.0 mL/minute; detector @ 220 nm; 25°C) showed one major peak, with a purity of 99.8%.

#### Preparation of Example 4




Example 4 was prepared from 6-bromo-DL-tryptophan ethyl ester hydrochloride, i.e., Intermediate 9, as depicted in the following synthetic scheme. Intermediate 9 was prepared from 4-bromo-2-nitrotoluene in four well-known synthetic steps. 4-Bromo-2-nitrotoluene is a commercially available compound from Aldrich Chemical Co., Milwaukee, WI.




- 43 -


5



10




15




20


- 44 -



Intermediate 9





Intermediate 13



Intermediate 14


- 45 -

5




10

Intermediate 16



29%



Example 4

15

Intermediate 11Preparation of 6-Bromoindole

20

The synthesis of Intermediate 11 is disclosed in M.P. Moyer et al., *J. Org. Chem.*, 51, page 5106 (1986); and K.L. Rinehart et al., *J. Am. Chem. Soc.*, 109, page 3378 (1987).

A mixture of 4-bromo-2-nitrotoluene (20 g, 93 mmol), pyrrolidine (8 mL, 93 mmol), and N,N-dimethylformamide dimethyl acetal (37 mL, 278 mmol)

- 46 -

in anhydrous N,N-dimethylformamide (200 mL) was heated at 110°C under a nitrogen blanket for 2 hours. The mixture was cooled, then diluted with water (1.5 L) and extracted with ether (3 x 500 mL).  
5 The combined organic extracts were washed with brine (500 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, and filtered. The solvent was removed under reduced pressure to provide Intermediate 10 as dark purple crystals which were suitable for use without further purification (27.9  
10 g): TLC R<sub>f</sub> (1:4 ethyl acetate/chloroform)=0.77.

Zinc granules (30 g, 459 mmol) were added over 30 minutes to a solution of crude Intermediate 10 (25 g, 93 mmol) in acetic acid (400 mL) and water (100 mL) at 75°C, after which the mixture was  
15 stirred for an additional 1 hour. The suspension was cooled, filtered, and extracted with ethyl acetate (2 x 2 L). The combined organic extracts were washed with water (2 x 2 L), saturated NaHCO<sub>3</sub> solution (2 x 500 mL), and brine (2 x 500 mL), dried  
20 over magnesium sulfate (MgSO<sub>4</sub>), and filtered. The solvent was removed under reduced pressure to yield a purple residue which was purified by flash column chromatography, eluting with hexanes/methylene chloride (1:1), to provide 6-bromoindole (Intermediate  
25 11) as a blue-white powder (4.9 g, 27% over two steps): TLC R<sub>f</sub> (1:9 ethyl acetate/chloroform)=0.81. The proton NMR spectrum (300 MHz, CDCl<sub>3</sub>) was identical to the known compound.

- 47 -

Intermediate 9Preparation of 6-Bromo-DL-tryptophan  
ethyl ester hydrochloride

5           The synthesis of Intermediate 12 is disclosed in T.L. Gilchrist et al., *J. Chem. Soc., Chem. Commun.*, page 1089 (1979); U. Schmidt et al., *Liebigs Ann. Chem.*, page 785 (1985); and U. Schmidt  
10 et al., *Tetrahedron Lett.*, 23, page 4911 (1982).

          Ethyl bromopyruvate (10 mL, 80 mmol) was added dropwise to a biphasic mixture of hydroxylamine sulfate (6.6 g, 40 mmol) in water (50 mL) and chloroform (50 mL), and the mixture was stirred at  
15 room temperature for 3 hours. The mixture was filtered, and the organic phase was concentrated under reduced pressure to yield the oxime Intermediate 15 as a white solid, which was used without further  
purification (15.9 g, 95%): TLC  $R_f$  (1:4 ethyl acetate/chloroform)=0.12.  
20

          Powdered sodium carbonate (4.0 g, 37 mmol) was added to a dark green solution of Intermediate 11 (4.9 g, 25 mmol) and Intermediate 15 (7.9 g, 37 mmol) in methylene chloride (50 mL) at room temperature under a nitrogen blanket. The mixture was  
25 stirred for 7 hours, after which the mixture was diluted with ethyl acetate (150 mL), washed with brine (100 mL), dried over  $MgSO_4$ , and filtered. The solvents were removed under reduced pressure to  
provide Intermediate 12 as a dark green oil which  
30 was suitable for use without further purification (9.6 g): TLC  $R_f$  (1:9 ethyl acetate/chloroform)=0.18.

          Aluminum amalgam was prepared by sequentially dipping several pieces of aluminum foil (5 g



- 48 -

total) in 1 N sodium hydroxide, saturated mercury (II) chloride solution, water, and ethanol. The aluminum pieces then were added to a solution of crude Intermediate 12 in ether (100 mL) and water (5 mL) at room temperature. The gray suspension was stirred for 24 hours, then filtered. The organic layer was washed with brine (100 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, and filtered. The solvents were removed under pressure to provide a dark green residue, which was treated with a ethereal HCl (1 M in ether). The solid was collected by vacuum filtration, and dried in a vacuum oven at 60°C overnight to provide Intermediate 9 as a dark red powder (3.3 g, 38% over two steps): <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 11.21 (s, 1H), 8.40 (s, 2H), 7.57 (s, 1H), 7.48 (d, J=8.4 Hz, 1H), 7.27 (s, 1H), 7.16 (d, J=7.0 Hz, 1H), 4.24-4.18 (m, 3H), 4.09 (q, J=6.8 Hz, 2H), 1.10 (t, J=7.3 Hz, 3H).

### Intermediate 13

#### Preparation of a *cis*-β-carboline

A solution of saturated aqueous sodium carbonate (50 mL) was added to a dark green solution of Intermediate 9 (3.4 g, 10 mmol) in methylene chloride (100 mL), and the resulting mixture was shaken for 10 minutes. The organic layer then was dried over Na<sub>2</sub>SO<sub>4</sub> and filtered. The solution was concentrated under reduced pressure to a volume of 20 mL and cooled to 0°C under a nitrogen blanket. Piperonal (1.7 g, 12 mmol) was added to the solution, followed by the addition of trifluoroacetic acid (1.5 mL, 20 mmol), after which the solution was

- 49 -

warmed to room temperature overnight. The reaction mixture then was diluted with ethyl acetate, and neutralized with aqueous  $\text{NaHCO}_3$  (100 mL). The organic layer was washed with brine (50 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and the solvent was removed under reduced pressure to provide a brown oil. This residue was purified by flash column chromatography, eluting with ethyl acetate/chloroform (1:19), to provide Intermediate 13 as a light yellow solid (0.88 g, 20%): TLC  $R_f$  (1:9 ethyl acetate/chloroform)=0.60;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  10.48 (s, 1H), 7.42 (s, 1H), 7.37 (d,  $J=10.0$  Hz, 1H), 7.09 (d,  $J=9.9$  Hz, 1H), 6.90 (d,  $J=6.2$  Hz, 2H), 6.83 (s, 1H), 6.00 (s, 2H), 5.15 (s, 1H), 4.16 (q,  $J=6.3$  Hz, 2H), 3.85-3.80 (m, 1H), 2.90-2.80 (m, 1H), 1.24 (t,  $J=6.9$  Hz, 3H). The *trans*- $\beta$ -carboline also was obtained as a light yellow solid, but was not characterized (0.60 g, 14%): TLC  $R_f$  (1:9 ethyl acetate/chloroform)=0.33.

#### Intermediate 14

#### Preparation of a *cis*-2-chloroacetyl- $\beta$ -carboline

Chloroacetyl chloride (0.2 mL, 2.6 mmol) was added dropwise to a solution of Intermediate 13 (0.88 g, 2.0 mmol) and triethylamine (0.4 mL, 2.6 mmol) in methylene chloride (10 mL) at  $0^\circ\text{C}$  under a nitrogen blanket. The mixture was slowly warmed to room temperature and stirred for 5 hours. The resulting white suspension was diluted with methylene chloride (100 mL), washed with brine (100 mL), dried over  $\text{Na}_2\text{SO}_4$ , and filtered. The solvent was removed under reduced pressure to provide Intermediate 14 as

- 50 -

an amber foam, which was used without further purification (1.03 g): TLC  $R_f$  (1:9 ethyl acetate/chloroform)=0.76.

5     Intermediate 16

A mixture of crude Intermediate 14 (1.0 g, 1.9 mmol) and methylamine (4 mL, 7.7 mmol, 2.0 M in THF) in methanol (10 mL) was heated at reflux under a nitrogen blanket for 2 hours, after which the resulting orange suspension was cooled to room temperature. The solids were collected by vacuum filtration and dried in a vacuum oven at 60°C overnight to provide Intermediate 15 as a white powder (0.45 g, 49% over two steps): mp 324-330°C; TLC  $R_f$  (1:9 methanol/chloroform)=0.78;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  11.18 (s, 1H), 7.53 (d,  $J$ =8.4 Hz, 1H), 7.49 (s, 1H), 7.14 (d,  $J$ =8.3 Hz, 1H), 6.86 (s, 1H), 6.78 (s, 2H), 6.11 (s, 1H), 5.93 (s, 2H), 4.39 (dd,  $J$ =4.4, 11.5 Hz, 1H), 4.17 (d,  $J$ =17.1 Hz, 1H), 3.95 (d,  $J$ =17.2 Hz, 1H), 3.52 (dd,  $J$ =4.5, 15.9 Hz, 1H), 3.01 (d,  $J$ =11.7 Hz, 1H), 2.93 (s, 3H); API MS  $m/z$  470  $[\text{C}_{22}\text{H}_{18}\text{BrN}_3\text{O}_4+\text{H}]^+$ . Anal. Calcd. for  $\text{C}_{22}\text{H}_{18}\text{BrN}_3\text{O}_4$ : C, 56.42; H, 3.87; N, 8.97. Found: C, 56.34; H, 3.92; N, 8.82. The stereochemistry of Intermediate 15 was confirmed to be the desired *cis* isomer by a series of NOE difference experiments: positive NOE enhancements from the C12a proton at 4.39 ppm to the C6 proton at 6.11 ppm and a C12 proton at 3.52 ppm; a positive NOE enhancement from the C6 proton at 6.11 ppm to the C12a proton at 4.39 ppm.

- 51 -

Example 4


Preparation of (+-, *cis*)-6-Benzo[1,3]dioxol-5-yl-  
2-methyl-9-phenyl-2,3,6,7,12,12a-hexahydro-  
pyrazino[1',2':1,6]pyrido[3,4-b]indole-1,4-dione

Phenyl boronic acid (0.09 g, 0.74 mmol)  
was added to a degassed mixture of Intermediate 16  
(0.172 g, 0.37 mmol), barium hydroxide octahydrate  
(0.58 g, 1.85 mmol), and bis(triphenylphosphine)-  
palladium (II) chloride (0.05 g, 0.07 mmol) in etha-  
nol (10 mL), and the resulting mixture was stirred  
at room temperature for 48 hours. An additional  
portion of phenyl boronic acid (0.09 g, 0.74 mmol)  
then was added, after which the mixture was stirred  
at room temperature for an additional 12 hours. The  
palladium catalyst was removed by vacuum filtration  
through a plug of Celite, and the resulting filtrate  
was concentrated under reduced pressure and purified  
by flash column chromatography, eluting with methy-  
lene chloride/ethyl acetate (4:1), to provide a  
white solid. This solid was further purified by  
recrystallization from methylene chloride followed  
by vacuum filtration to yield a solid, which was  
dried overnight under vacuum at 60°C to provide  
Example 4 as a white solid (0.049 g, 29%): mp 198-  
208°C; TLC R<sub>f</sub> (1:4 ethyl acetate/chloroform)=0.37; <sup>1</sup>H  
NMR (500 MHz, CDCl<sub>3</sub>): δ 7.91 (s, 1H), 7.46 (s, 1H),  
7.37 (d, J=8.4 Hz, 2H), 7.27-7.25 (m, 4H), 6.95 (s,  
1H), 6.79 (s, 1H), 6.72-6.67 (m, 2H), 5.93 (s, 2H),  
5.29 (s, 1H), 4.34 (dd, J=4.0, 4.11 Hz, 1H), 4.12  
(dd, J=1.2, 1.1 Hz, 1H), 3.51 (dd, J=4.2, 4.2 Hz,  
1H), 3.00 (s, 3H), 2.93-2.88 (m, 2H) ppm; API MS m/z  
467 [C<sub>28</sub>H<sub>23</sub>N<sub>3</sub>O<sub>4</sub>+H]<sup>+</sup>. HPLC analysis (Symmetry C18 Col-

- 52 -

umn, 150 x 3.5 mm. Retention Time=10.9 minutes;  
30/0.85:70/0.1 acetonitrile/TFA:water/TFA to  
100/0.85 acetonitrile/TFA over 20 minutes; flow=1.0  
mL/min; detector at 220 nm; 25°C) showed one peak,  
5 with a purity of 91.5%. Chiral HPLC analysis  
(Chiralcel OD Column, 250 x 4.6 mm, Retention  
Times=12.9 and 17.4 min; 1:1 isopropanol/hexanes;  
flow=0.5 mL/min; detector at 222 nm; 25°C) showed  
two major peaks, with a ratio of 54:42 and with a  
10 total purity of 96.6%. The relative stereochemistry  
of Intermediate 4 was confirmed to be the desired  
*cis* isomer by a series of NOE difference experi-  
ments: a positive NOE enhancement from the C12a  
proton at 4.34 ppm to the C6 proton at 6.79 ppm; a  
15 positive NOE enhancement from the C6 proton at 6.79  
ppm to the C12a proton at 4.34 ppm.

#### Preparation of Example 5




Example 5 was prepared from indole-6-  
30 carboxylic acid as depicted in the following syn-  
thetic scheme. Indole-6-carboxylic acid is a com-  
mercially available compound. Aspects of the fol-  
lowing synthetic scheme are disclosed in Y.


- 53 -

Yokoyama et al., *Tetrahedron Lett*, 40, p. 7803 (1990); and H.R. Snyder et al., *J. Am Chem. Soc.*, 77, p. 1257 (1955).

5




10



15

Intermediate 17


20



25

Intermediate 18


- 54 -




Intermediate 19



Intermediate 20




Intermediate 21




Intermediate 22

- 55 -

5





53% cis  
43% trans



Intermediate 23

10



Example 5

20

Intermediate 17Preparation of N-Acetyl-6-carboxytryptophan

25

A mixture of indole-6-carboxylic acid (5.0 g, 31 mmol), DL-serine (3.25 g, 31 mmol), and acetic anhydride (8.8 mL, 93 mmol) in glacial acetic acid (50 mL) was heated under a nitrogen blanket at 80°C for 24 hours. The resulting brown solution was cooled to room temperature, then the solvent was removed under reduced pressure to provide Intermediate 17 as a brown foam, which was used without fur-

30



- 56 -

ther purification (11.0 g):  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ ):  
 $\delta$  8.08 (s, 1H), 7.72 (s, 2H), 7.40 (s, 1H), 4.61-  
4.53 (m, 1H), 3.45-3.35 (m, 1H), 3.23-3.13 (m, 1H),  
1.92 (s, 3H) ppm.

5

### Intermediate 18

#### Preparation of 6-Carboxytryptophan

A solution of crude Intermediate 17 (11.0  
10 g) was heated in 15% aqueous sodium hydroxide for 18  
hours. The reaction mixture was cooled to  $0^\circ\text{C}$ , then  
acidified to pH 5 with acetic acid, after which the  
solution was concentrated under reduced pressure.  
The residue was slurried in water (500 mL), then the  
15 solids were collected by filtration under reduced  
pressure and dried in a vacuum oven at  $100^\circ\text{C}$  for 18  
hours to yield Intermediate 18 as a gray solid (2.6  
g, 34% over two steps). A second portion of Inter-  
mediate 18 was recovered from the filtrate (0.75 g,  
20 10% over two steps):  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$   
11.30 (s, 1H), 8.00 (s, 1H), 7.68-7.52 (m, 2H), 7.40  
(s, 1H), 3.50-3.40 (m, 1H), 3.39-3.25 (m, 1H), 3.10-  
2.92 (m, 1H) ppm.

### 25 Intermediates 19-21

#### Preparation of 6-Methylcarboxytryptophan

Thionyl chloride (0.84 mL, 11 mmol) was  
added dropwise to a suspension of Intermediate 18  
30 (0.75 g, 302 mmol) in methanol (15 mL) at  $0^\circ\text{C}$  under  
a nitrogen blanket, then the mixture was warmed to  
room temperature and stirred for 20 hours. The  
resulting solution was diluted with ethyl acetate

- 57 -

(200 mL), extracted with saturated sodium bicarbonate ( $\text{NaHCO}_3$ ) solution (2 x 50 mL) and acidified to pH 6 with acetic acid. The resulting slurry was filtered, then the solid was washed with several small portions of water, and dried in a vacuum oven at room temperature to yield a mixture of Intermediates 19 and 20 as a brown solid (400 mg, 51%).

Thionyl chloride (0.45 mL, 6.0 mmol) was added dropwise to a slurry of Intermediates 19 and 20 (316 mg, 1.21 mmol) in methanol (15 mL) at 0°C under a nitrogen blanket. This mixture was warmed to room temperature and stirred for 3 hours, then the mixture was heated to 50°C to stir for an additional 2 hours. The solvent was removed under reduced pressure, and the residue was diluted with ethyl acetate (200 mL). The solution was washed with saturated  $\text{NaHCO}_3$  (20 mL), water (20 mL), and brine (20 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and the solvent was removed under reduced pressure to yield Intermediate 21 as a brown oil (310 mg, 93%): TLC  $R_f$  (5:1 methylene chloride/ethyl acetate)=0.1:  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.11 (bs, 1H), 8.07 (s, 1H), 7.80 (d,  $J=8.0$  Hz, 1H), 7.61 (d,  $J=8.0$  Hz, 1H), 7.19 (s, 1H), 3.93 (s, 3H), 3.90-3.82 (m, 1H), 3.71 (s, 1H), 3.31-3.22 (m, 1H), 3.13-3.05 (m, 1H) ppm.

### Intermediate 22

#### Preparation of $\beta$ -Carboline

A mixture of Intermediate 21 (495 mg, 1.79 mmol), piperonal (479 mg, 3.2 mmol), and p-toluene sulfonic acid monohydrate (80 mg) was refluxed in toluene with water removal via Dean-Stark trap for 5

- 58 -

hours under a nitrogen blanket. The dark brown mixture was cooled to room temperature and diluted with ethyl acetate (250 mL). The mixture then was washed successively with saturated  $\text{NaHCO}_3$  (30 mL), water (20 mL), and brine (20 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered, and the solvent was removed under reduced pressure. The residue was slurried in a mixture of methylene chloride/ethyl acetate (5:1, 10 mL) and the slurry was filtered under pressure to yield a mixture of the cis- and trans- Intermediate 22, which were used without characterization (311 mg, 42%).

### Intermediate 23

#### Preparation of cis-2-Chloroacetyl- $\beta$ -carboline

Chloroacetyl chloride (0.076 mL, 0.99 mmol) was added dropwise to a mixture of Intermediate 22 (311 mg, 0.76 mmol, mixture of cis and trans isomers) and triethylamine (0.138 mL, 0.99 mmol) in chloroform (10 mL) at 0°C under a nitrogen blanket. The resulting mixture was stirred at 0°C for 1 hour, then warmed to room temperature and stirred for an additional 2 hours. The solution was diluted with ethyl acetate (100 mL), washed with saturated  $\text{NaHCO}_3$  solution (10 mL), water (10 mL), and brine (10 mL), dried over sodium sulfate ( $\text{Na}_2\text{SO}_4$ ), and the solvent was removed under reduced pressure to provide a yellow foam. The mixture was purified by flash column chromatography, eluting with hexanes/ethyl acetate (2:1), to yield Intermediate 23 as a white solid, which was not characterized (195 mg, 53%): TLC  $R_f$  (2:1 hexanes/ethyl acetate)=0.62. The latter

- 59 -

eluting trans-2-chloroacetyl- $\beta$ -carboline was obtained as a white solid which also was not characterized (160 mg, 43%): TLC  $R_f$  (2:1 hexanes/ethyl acetate)=0.40.

#### Example 5

Preparation of (+-, cis)-6-Benzo[1,3]dioxol-5-yl-2-methyl-1,4-dioxo-1,2,3,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indole-9-carboxylic acid methyl ester


A mixture of Intermediate 23 (132 mg, 0.27 mmol) and methylamine (6.2 mL, 12.4 mmol, 2 M solution in THF) in methylene chloride (18 mL) was refluxed under a nitrogen blanket for 4 hours. The resulting solution was concentrated under reduced pressure to yield a yellow solid, which was stirred in methanol (4 mL) for 1 hour. The solid was isolated by filtration under reduced pressure, washed with methanol (5 x 1 mL), and dried in a vacuum oven at 70°C for 17 hours to provide Example 5 as an off-white solid (117 mg, 96%): mp 294-295°C; TLC  $R_f$  (5:1:0.5 methylene chloride/ethyl acetate/methanol) =0.65;  $^1H$  NMR (300 MHz,  $CDCl_3$ ):  $\delta$  8.25 (bs, 1H), 8.03 (s, 1H), 7.84 (d,  $J$ =8.7 Hz, 1H), 7.60 (d,  $J$ =8.7 Hz, 1H), 6.86 (dd,  $J$ =8.0, 1.7 Hz, 1H), 6.73 (d,  $J$ =1.7 Hz, 1H), 6.69 (d,  $J$ =8.0 Hz, 1H), 6.20 (s, 1H), 5.88 (s, 1H), 5.86 (s, 1H), 4.30 (dd,  $J$ =11.6, 4.5 Hz, 1H), 4.11 (dd,  $J$ =17.6, 1.3 Hz, 1H), 4.00-3.88 (m, 4H), 3.78 (dd,  $J$ =16.1, 4.5 Hz, 1H), 3.27-3.16 (m, 1H), 3.05 (s, 3H) ppm; CI MS  $m/z$  448  $[C_{24}H_{21}N_3O_6+H]^+$ , Anal. Calcd. for  $C_{24}H_{21}N_3O_6$ : C, 64.42; H, 4.73; N, 9.39. Found: C, 64.12; H, 4.61; N, 9.12. HPLC analysis (Aquasil C18 Column, 100 x 4.6

- 60 -

mm, Retention Time=10.3 minutes; 45:55/0.03 acetonitrile:water/TFA; flow=0.50 mL/min.; detector @ 254 nm; temperature ambient) showed one peak, with a purity of 99.7%. The stereochemistry of Example 5 was confirmed to be the cis isomer by a series of NOE difference experiments: a positive NOE enhancement from the C12a proton at 4.30 ppm to the C6 proton at 6.20 ppm; a positive NOE enhancement from the C6 proton at 6.20 ppm to the C12a proton at 4.30 ppm.

#### Preparation of Example 6

(+,-,trans)-6-Benzo[1,3]dioxol-5-yl-2-methyl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indole-9-carboxylic acid




Example 6 was prepared from Example 5 by the following procedure.

A mixture of Example 5 (0.125 g, 0.28 mmol), 2 N sodium hydroxide solution (2 mL), THF (6 mL), and methanol (2 mL) was heated at 80°C for 24 hours. The resulting yellow solution was cooled to room temperature and diluted with water, then the




- 61 -

solution was acidified to pH 6 with acetic acid. The resulting solution was concentrated under reduced pressure to a volume of 20 mL, and was allowed to stand overnight at room temperature. The resulting solid precipitate was isolated by filtration under reduced pressure, then further purified by stirring in diethyl ether (4 mL) at 40°C for 2 hours. The resulting solids were isolated by filtration under reduced pressure, washed with diethyl ether (5 x 1 mL), and dried in a vacuum oven at 90°C for 4 hours to provide Example 6 as an off-white solid (0.093, 78%): mp 246-252°C; <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD): δ 8.05 (s, 1H), 7.76 (d, J=8.6 Hz, 1H), 7.56 (d, J=8.14 Hz, 1H), 6.99 (s, 1H), 6.83-6.73 (m, 2H), 6.66 (dd, J=8.0 Hz, 1H), 5.93 (s, 2H), 4.34-4.23 (m, 2H), 4.08 (d, J=17.3 Hz, 1H), 3.44 (dd, J=15.4, 4.2 Hz, 1H), 3.08-2.95 (m, 4H) ppm; ESI MS m/z 432 [C<sub>23</sub>H<sub>19</sub>N<sub>3</sub>O<sub>6</sub>-H]<sup>+</sup>. Anal. Calcd. for C<sub>23</sub>H<sub>19</sub>N<sub>3</sub>O<sub>6</sub>: C, 63.74; H, 4.42; N, 9.70. Found: C, 62.77; H, 4.61; N, 9.19. HPLC analysis (Aquasil C18 Column, 100 x 4.6 mm, Retention Time=25.4 minutes; 30:70/0.03 acetonitrile:water/TFA; flow=0.50 mL/min; detector at 254 nm; temperature ambient) showed one peak, with a purity of 98.9%. The stereochemistry of Example 6 was confirmed to be the trans isomer by a series of NOE difference experiments: no NOE enhancement from the C12a proton at 4.30 ppm to the C6 proton at 6.99 ppm; no NOE enhancement from the C6 proton at 6.99 ppm to the C12a proton at 4.30 ppm.


- 62 -

Preparation of Example 7

Example 7 was prepared from 6-cyanoindole as depicted in the following synthetic scheme. 6-Cyanoindole is a commercially available compound.




- 63 -



Intermediate 26

1. NaOH (1 eq), MeOH  
2. HOAc  
3. CH<sub>2</sub>Cl<sub>2</sub>, MeOH, reflux

46%



Intermediate 27


xylenes, p-TsOH

145°C

Dean-Stark

28% cis

23% trans




Intermediate 28


ClCOCH<sub>2</sub>Cl  
Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>



- 64 -



Intermediate 29



Example 7

MeNH<sub>2</sub>, THF  
→  
MeOH, 50°C  
44% over  
two steps

#### Intermediate 24

#### Preparation of Gramine

Acetic acid (6 ml) was added dropwise to aqueous dimethylamine (7.0 ml, 55.7 mmol) at a rate that maintained the solution below 5°C. To this mixture was added aqueous formaldehyde (4.2 ml, 55.7 mmol) dropwise. The resulting mixture then was added to a solution of 6-cyanoindole (6.6 g, 46.4

- 65 -

mmol) in acetic acid (30 ml) over 15 minutes at room temperature under a nitrogen blanket. The resulting dark yellow solution was stirred at room temperature for 3 hours, followed by dilution with 2 N sodium hydroxide (30 ml), then stored at 0°C for 12 hours. The solution was decanted, and the solvent was removed under reduced pressure to produce a white foam. The white foam was dissolved in water (200 mL), and adjusted to a basic pH with a saturated NaHCO<sub>3</sub> solution. The resulting mixture was extracted with ethyl acetate (3 x 200 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, and the solvent was removed under reduced pressure to yield Intermediate 24 as an off-white solid (6.24 g, 65%): <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 11.52 (s, 1H), 7.85 (s, 1H), 7.77 (d, J=8.2 Hz, 1H), 7.54 (d, J=2.4 Hz, 1H), 7.31 (dd, J=1.5, 8.3 Hz, 1H), 3.56 (s, 2H), 2.13 (s, 6H) ppm.

### Intermediate 25

#### Preparation of Diester

To a solution of diethyl formamidomalonate (6.61 g, 32.5 mmol) in ethanol (40 mL) under an argon blanket was added sodium ethoxide (12.0 mL, 32.5 mmol, 2.7 M solution in ethanol). The resulting mixture was stirred at room temperature for 30 minutes to yield a slurry. To the slurry was added Intermediate 24 (5.39 g, 27.1 mmol), followed by dimethyl sulfate (5.1 mL, 54.2 mmol). The resulting clear brown solution was stirred at room temperature, and after 15 minutes a precipitate formed. The resulting slurry was stirred at room temperature for 16 hours. Then the solids were removed by fil-

- 66 -

tration under reduced pressure, and the filtrate concentrated to yield diethyl ester Intermediate 25 as an orange solid which was used without further purification (10.9 g): TLC  $R_f$  (4:1 methylene chloride/ethyl acetate)=0.50.

### Intermediate 26

#### Preparation of Amino Diester

To a solution of Intermediate 25 (27.1 mmol) in methanol (120 mL) was added hydrochloric acid in diethyl ether (54 mL, 54.2 mmol, 1 M solution in diethyl ether). The resulting solution was allowed to sit overnight under an argon blanket. The mixture then was concentrated under reduced pressure, diluted with ethyl acetate (100 mL), and extracted with 2 N HCl (3 x 200 mL). The combined aqueous extracts were adjusted to pH 8 with  $\text{Na}_2\text{CO}_3$ , followed by extraction with methylene chloride (3 x 200 mL). The combined organic extracts were dried over  $\text{Na}_2\text{SO}_4$ , and the solvent was removed under reduced pressure to provide Intermediate 26 as an orange oil (2.57 g, 30%):  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.79 (bs, 1H), 7.69-7.64 (m, 2H), 7.34-7.30 (m, 2H), 4.27-4.15 (m, 4H), 3.48 (s, 2H), 1.99 (bs, 2H), 1.31-1.23 (m, 6H) ppm.

- 67 -

Intermediate 27Preparation of (+/-)-6-Cyanotryptophan Ethyl Ester

To a mixture of Intermediate 26 (2.57 g, 7.80 mmol) in ethanol (80 mL) was added a solution of sodium hydroxide (0.37 g, 9.25 mmol) in water (5 mL). The resulting orange solution was stirred at room temperature for 16 hours, after which an additional 0.5 equivalents of sodium hydroxide in water was added. The mixture was stirred for 5 hours, then acidified to pH 6 with acetic acid. Concentration of the solution under reduced pressure provided an oil which was dissolved in 2:1 methylene chloride/methanol (60/20 mL), then heated at reflux for 5 hours. The resulting slurry was cooled, then the solids were removed by filtration under reduced pressure. The filtrate was concentrated, dissolved in water (100 mL), then extracted with methylene chloride (3 x 100 mL). The organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, and the solvent was removed under reduced pressure to provide Intermediate 27 as an orange oil (0.91 g, 46%): <sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>): δ 11.40 (s, 1H), 7.81 (s, 1H), 7.68 (d, J=10.0 Hz, 1H), 7.45 (s, 1H), 7.30 (d, J=10.0 Hz, 1H), 4.00-3.96 (m, 2H), 3.60-3.57 (m, 1H), 3.03-2.94 (m, 2H), 2.08 (bs, 2H), 1.10-1.07 (m, 3H) ppm.

Intermediate 28Preparation of (+/-)-cis-β-Carboline

A solution of Intermediate 27 (0.49 g, 1.9 mmol), piperonal (0.37 g, 2.5 mmol) and p-toluene-sulfonic acid monohydrate (0.050 g, 0.26 mmol) in

- 68 -

xylenes (30 mL) was heated at 145°C under an argon blanket for 16 hours with water removal via Dean-Stark trap. The resulting orange solution was cooled, diluted with methylene chloride (100 mL), then washed with saturated NaHCO<sub>3</sub> (20 mL). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, then the solvent was removed under reduced pressure to provide an orange oil. The residue was purified by flash column chromatography, eluting with methylene chloride/ethyl acetate (6:1) to provide Intermediate 28 as orange solid (0.20 g, 27%): TLC R<sub>f</sub> (5:1 methylene chloride/ethyl acetate)=0.59; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 10.96 (s, H), 7.63-7.59 (m, 2H), 7.30 (d, J=9.9 Hz, 1H), 6.94-6.84 (m, 3H), 6.01 (s, 2H), 5.20 (d, J=9.9 Hz, 1H), 4.19 (q, J=7.3 Hz, 2H), 3.87-3.82 (m, 1H), 3.12-3.03 (m, 1H), 2.89-2.78 (m, 1H), 1.28 (t, J=7.1 Hz, 3H) ppm. The latter eluting trans isomer also was isolated as an orange solid (0.17 g, 23%): TLC R<sub>f</sub> (5:1 methylene chloride/ethyl acetate)=0.28; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 11.21 (s, 1H), 7.69 (s, 1H), 7.63 (d, J=9.9 Hz, 1H), 7.30 (d, J=9.9 Hz, 1H), 6.88-6.82 (m, 2H), 6.68 (d, J=9.0 Hz, 1H), 5.98 (s, 2H), 5.31 (s, 1H), 4.12-4.05 (m, 2H), 3.84-3.79 (m, 1H), 3.14-3.03 (dd, J=5.0, 15.0 Hz, 1H), 2.95-2.88 (dd, J=8.0, 15.0 Hz, 1H), 1.18 (t, J=7.5 Hz, 3H) ppm.

### Intermediate 29

#### Preparation of (+/-)-cis-2-Chloroacetyl-β-Carboline

Chloroacetyl chloride (0.055 mL, 69 mmol) was added to a solution of Intermediate 28 (0.20 g, 0.51 mmol) and triethylamine (0.10 mL, 0.72 mmol) in

- 69 -

methylen chloride (10 mL) at 0°C under an argon blanket, after which the solution was warmed to room temperature over 2 hours. The yellow solution then was diluted with methylen chloride (40 mL); washed with water (20 mL), and saturated NaHCO<sub>3</sub> solution (20 mL). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, and the solvent was removed under reduced pressure to afford Intermediate 29 as a yellow oil which was used without further purification: TLC R<sub>f</sub> (8:1 methylen chloride/ethyl acetate)=0.76.

#### Example 7

Preparation of (+-, cis)-6-Benzo[1,3]dioxol-5-yl-2-methyl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydro-pyrazino[1',2':1,6]pyrido[3,4-b]indole-9-carbonitrile


A solution of cis-2-chloroacetyl-β-carboline 7 (0.51 mmol) and methylamine (5.1 mL, 10.2 mmol, 2 M solution in THF) in methanol (9.0 mL) was heated at 50°C for 16 hours. The resulting solids were isolated by filtration under reduced pressure to provide Example 7 as a pale yellow solid (94 mg, 44%): mp 329-331°C; TLC R<sub>f</sub> (5:1:0.5 methylene chloride/ethyl acetate/methanol)=0.50; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 11.65 (s, H), 7.81 (s, 1H), 7.76 (d, J=8.2 Hz, 1H), 7.33 (d, J=8.4 Hz, 1H), 6.88 (s, 1H), 6.79 (s, 2H), 6.17 (s, 2H), 5.93 (s, 2H), 4.43-4.38 (dd, J=4.1, 11.5 Hz, 1H), 4.17 (d, J=17.3 Hz, 1H), 3.95 (d, J=17.3 Hz, 1H), 3.59 (dd, J=4.4, 16.0 Hz, 1H), 3.00 (dd, J=11.5, 15.7 Hz, 1H), 2.93 (s, 3H) ppm; ESI MS m/z 413 [C<sub>23</sub>H<sub>18</sub>N<sub>4</sub>O<sub>4</sub>-H]<sup>+</sup>. Anal. Calcd. for C<sub>23</sub>H<sub>18</sub>N<sub>4</sub>O<sub>4</sub>; C, 66.66; H, 4.38; N, 13.52. Found: C, 66.49; H, 4.43; N, 13.30. HPLC analysis

- 70 -

(Aquasil C18 Column, 100 x 4.6 mm, Retention Time=9.74 min; 45:55 0.03% acetonitrile:water/TFA; flow=0.50 mL/min; detector @ 254 nm; temperature ambient) showed one peak, with a purity of 100.0%.


The stereochemistry of analog Example 7 was confirmed to be the desired cis isomer by a series of NOE difference experiments: a positive NOE enhancement from the C12a proton at 4.40 ppm to the C6 proton at 6.17 ppm; a positive NOE enhancement from the C6 proton at 6.17 ppm to the C12a proton at 4.40 ppm.

#### Preparation of Example 8




Example 8 was prepared from 3-methyl-2-nitrophenol as depicted in the following synthetic scheme. Intermediate 31 was prepared as disclosed in G.M. Carrera et al., *Synlet*, p. 93 (1994) and M.P. Moyer et al., *J. Org. Chem.*, 51, p. 5106 (1986), and converted to Intermediate 32 as disclosed in Y. Yokohama, *Tetrahedron Lett.*, 40, p. 7803 (1999) and H.R. Snyder et al., *J. Am. Chem. Soc.*, 77, p. 1257 (1955).


- 71 -



5




10



Intermediate 31

15



Intermediate 32



20

25





- 72 -



5



10




20



25

30

- 73 -



Example 8

### Intermediate 31

#### Preparation of 7-Benzzyloxyindole

Benzyl bromide (15.4 ml, 130 mmol) was added dropwise to a solution of 3-methyl-2-nitrophenol (20.0 g, 130 mmol) and sodium hydroxide (6.0 g, 144 mmol) in anhydrous N,N-dimethylformamide (100 mL) at 0°C under a nitrogen blanket. The resulting red solution was slowly warmed to room temperature, stirring for a total of 17 hours. The mixture then was diluted with ethyl acetate (700 ml), stirred for 15 minutes, and the solids were removed by filtration under reduced pressure. The bright yellow filtrate was concentrated under reduced pressure to provide Intermediate 30 as a dark yellow residue, which was used without further purification (31.8 g): TLC R<sub>f</sub> (5:1 hexanes/ethyl acetate)=0.50.

A mixture of Intermediate 30 (31.8 g, 130 mmol), pyrrolidine (11.0 mL, 130 mmol), and N,N-dimethylformamide dimethyl acetal (52 mL, 392 mmol) in anhydrous N,N-dimethylformamide (100 mL) was

- 74 -

heated at 110°C under a nitrogen blanket for 3 hours. The cooled mixture was diluted with water (1.5 L), extracted with ether (2 x 500 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, and filtered. The solvent was removed under reduced pressure to afford the intermediate enamine as a dark red oil, which was immediately used in the next step without purification.

A solution of the enamine in methanol (150 mL) and acetic acid (5 mL) was treated with a catalytic amount of 10% palladium on carbon (2 g, 10% wet), and the resulting mixture was stirred under a hydrogen atmosphere (50 psi) at room temperature for 2 hours. The palladium catalyst was removed by vacuum filtration through MgSO<sub>4</sub> (10 g), and the filtrate was concentrated under reduced pressure to provide a dark brown residue. The residue was purified by flash column chromatography, eluting with ethyl acetate/chloroform (1:19), to provide Intermediate 31 as an off-white solid (3.82 g, 21% over three steps): TLC R<sub>f</sub> (4:1 hexanes/ethyl acetate)=0.61; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 11.18 (s, 1H), 7.56 (d, J=7.2 Hz, 2H), 7.43-7.38 (m, 3H), 7.35 (s, 1H), 7.23 (d, J=2.9 Hz, 1H), 6.88 (t, J=7.6 Hz, 1H), 6.72 (d, J=7.6 Hz, 1H), 6.39 (t, J=2.5 Hz, 1H), 5.26 (s, 2H) ppm.

### Intermediate 32

#### Preparation of (+/-)-7-Benzoyloxytryptophan

Acetic anhydride (0.5 mL, 5.2 mmol) was added to a slurry of Intermediate 31 (0.575 g, 2.6 mmol) and DL-serine (0.270 g, 2.6 mmol) in anhydrous acetic acid (5 mL) at room temperature under a ni-

- 75 -

trogen blanket. The resulting orange solution was heated to reflux and stirred for 24 hours. The cooled mixture then was concentrated under reduced pressure to provide the intermediate N-acetyltryptophan as dark red oil, which was used immediately without purification.

A suspension of the intermediate N-acetyltryptophan in 3 N sodium hydroxide was heated at reflux for 17 hours, then cooled to room temperature. The mixture was diluted with water (50 mL), washed with diethyl ether (50 mL), slurried with charcoal (1 g), and filtered through a plug of silica gel (20 g), eluting with water (100 mL). The filtrate was acidified to pH 4 with 6 N HCl (1 mL), and the solution was filtered through a plug of DOWEX 50X8-100 ion-exchange resin (100 g), eluting with concentrated ammonium hydroxide/methanol (1:3), to provide Intermediate 32 as a gray powder (0.040 g, 5% over two steps):  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.51 (d,  $J=7.0$  Hz, 2H), 7.39-7.29 (m, 3H), 7.17 (d,  $J=7.9$  Hz, 1H), 7.04 (s, 1H), 6.91 (t,  $J=7.6$  Hz, 1H), 6.68 (d,  $J=7.7$  Hz, 1H), 5.21 (s, 2H), 4.72-4.68 (m, 1H), 3.34-3.29 (m, 1H), 3.16-3.09 (m, 1H) ppm.

### Intermediate 33

#### Preparation of (+/-)-7-Benzoyloxytryptophan Methyl Ester Hydrochloride

Thionyl chloride (0.4 mL, 4.8 mmol) was added to a suspension of Intermediate 32 (0.600 g, 1.9 mmol) in anhydrous methanol (10 mL) at 0°C under a nitrogen blanket. The resulting mixture was heated to 50°C and stirred for 1 hour, then cooled

- 76 -

to room temperature. The resulting dark green solution was concentrated under reduced pressure to provide Intermediate 33 as a light green powder which was used without further purification (0.710 g, 100%): TLC  $R_f$  (4:1 chloroform/methanol)=0.87;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  11.18 (s, 1H), 8.48 (bs, 2H), 7.55 (d,  $J=7.2$  Hz, 1H), 7.43-7.41 (m, 3H), 7.38 (s, 1H), 7.34 (d,  $J=7.2$  Hz, 1H), 7.10-6.91 (m, 1H), 6.75 (d,  $J=7.6$  Hz, 1H), 5.26 (s, 2H), 4.42 (s, 1H), 3.66 (s, 3H), 3.35-3.26 (m, 2H) ppm.

#### Intermediate 34

#### Preparation of (+/-)-cis- $\beta$ -Carboline

A solution of Intermediate 33 (0.710 g, 2.0 mmol) and piperonal (0.300 g, 2.0 mmol) in anhydrous isopropanol (10 mL) was heated at reflux under a nitrogen blanket for 6 hours. The resulting orange solution was cooled to room temperature, neutralized with saturated  $\text{NaHCO}_3$  solution (1 mL), then the solvent was removed under reduced pressure to provide brown solid. The residue was purified by flash column chromatography, eluting with chloroform/ethyl acetate (19:1), to provide Intermediate 34 as a colorless oil (0.120 g, 13%): TLC  $R_f$  (4:1 chloroform/ethyl acetate)=0.57;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.70 (s, 1H), 7.44-7.26 (m, 5H), 7.17 (d,  $J=7.8$  Hz, 1H), 7.04 (t,  $J=7.7$  Hz, 1H), 6.88-6.70 (m, 4H), 5.94 (s, 2H), 5.16 (s, 2H), 3.93 (dd,  $J=4.1$ , 11.0 Hz, 1H), 3.81 (s, 3H), 3.22-3.17 (m, 1H), 3.03-2.95 (m, 1H) ppm. The latter eluting trans isomer was also isolated as a colorless oil, but was not

- 77 -

characterized (0.190 g, 21%): TLC  $R_f$  (4:1 chloroform/ethyl acetate)=0.51.

### Intermediate 35

#### 5 Preparation of (+/-)-cis-2-Chloroacetyl- $\beta$ -carboline

Chloroacetyl chloride (0.03 mL, 0.3 mmol) was added to a solution of Intermediate 34 (0.120 g, 0.3 mmol) and triethylamine (0.05 mL, 0.3 mmol) in methylene chloride (5 mL) at 0°C under an argon blanket, then warmed to room temperature over 1 hour. The solvent then was removed under reduced pressure to yield Intermediate 35 as a yellow powder which was used without further purification: TLC  $R_f$  (4:1 chloroform/ethyl acetate)=0.71.

### Example 8


#### 20 Preparation of (+-, cis)-6-Benzo[1,3]dioxol-5-yl-8-benzyloxy-2-methyl-2,3,6,7,12,12a-hexahydropyrazino[1'2':1,6]pyrido[3,4-b]indole-1,4-dione

A solution of Intermediate 35 (0.3 mmol) and methylamine (0.7 mL, 1.2 mmol, 2 M solution in THF) in methanol (5 mL) was heated at 50°C for 20 hours. The resulting solids were isolated by filtration under reduced pressure to provide Example 8 as a pale yellow powder (0.092 g, 71% over two steps): mp 150-161°C; TLC  $R_f$  (4:1 chloroform/ethyl acetate)=0.19;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.25 (s, H), 7.45-7.34 (m, 6H), 7.26-7.22 (m, 1H), 7.06 (t, J=7.9 Hz, 1H), 6.84 (d, J=8.0 Hz, 1H), 6.74-6.65 (m, 2H), 6.17 (s, 1H), 5.86 (s, 2H), 5.14 (s, 2H), 4.23 (dd, J=4.3, 11.5 Hz, 1H), 3.95 (dd, J=17.4, 52.8 Hz,

- 78 -


2H), 3.75 (dd, J=4.5, 11.4 Hz, 1H), 3.19 (dd, J=4.3, 11.7 Hz, 1H), 3.01 (s, 3H) ppm; ESI MS m/z 496 [C<sub>29</sub>H<sub>25</sub>N<sub>3</sub>O<sub>5</sub>+H]<sup>+</sup>. Anal. Calcd. for C<sub>29</sub>H<sub>25</sub>N<sub>3</sub>O<sub>5</sub>: C, 70.29; H, 5.09; N, 8.48. Found: C, 69.71; H, 5.09; N, 8.41. The stereochemistry of Example 8 was confirmed to be the desired cis isomer by a series of NOE difference experiments: a positive NOE enhancement from the C12a proton at 4.23 ppm to the c6 proton at 6.17 ppm; a positive NOE enhancement from the C6 proton at 6.17 ppm to the C12a proton at 4.23 ppm.

The following compounds are additional examples of compounds of structural formula (I) that can be prepared by methods analogous to the preparation of Examples 1 through 8.




### Example 9

- 79 -




Example 10




Example 11




- 80 -



Example 12



Example 13



Example 14

- 81 -

Compounds of the present invention can be formulated into tablets for oral administration. For example, a compound of formula (I) can be formed into a dispersion with a polymeric carrier by the coprecipitation method set forth in WO 96/38131, incorporated herein by reference. The coprecipitated dispersion can be blended with excipients, then pressed into tablets, which optionally are film-coated.

The compounds of structural formula (I) were tested for an ability to inhibit PDE5. The ability of a compound to inhibit PDE5 activity is related to the  $IC_{50}$  value for the compound, i.e., the concentration of inhibitor required for 50% inhibition of enzyme activity. The  $IC_{50}$  value for compounds of structural formula (I) were determined using recombinant human PDE5.

The compounds of the present invention typically exhibit an  $IC_{50}$  value against recombinant human PDE5 of less than about 50  $\mu M$ , and preferably less than about 25  $\mu M$ , and more preferably less than about 15  $\mu M$ . The compounds of the present invention typically exhibit an  $IC_{50}$  value against recombinant human PDE5 of less than about 1  $\mu M$ , and often less than about 0.05  $\mu M$ . To achieve the full advantage of the present invention, a present PDE5 inhibitor has an  $IC_{50}$  of about 0.1 nM to about 15  $\mu M$ .

The production of recombinant human PDEs and the  $IC_{50}$  determinations can be accomplished by well-known methods in the art. Exemplary methods are described as follows:

- 82 -

**EXPRESSION OF HUMAN PDEs****Expression in *Saccharomyces cerevisiae* (Yeast)**

5                   Recombinant production of human PDE1B,  
PDE2, PDE4A, PDE4B, PDE4C, PDE4D, PDE5, and PDE7 was  
carried out similarly to that described in Example 7  
of U.S. Patent No. 5,702,936, incorporated herein by  
reference, except that the yeast transformation  
10       vector employed, which is derived from the basic  
ADH2 plasmid described in Price et al., *Methods in  
Enzymology*, 185, pp. 308-318 (1990), incorporated  
yeast ADH2 promoter and terminator sequences and the  
*Saccharomyces cerevisiae* host was the protease-defi-  
15       cient strain BJ2-54 deposited on August 31, 1998  
with the American Type Culture Collection, Manassas,  
Virginia, under accession number ATCC 74465. Trans-  
formed host cells were grown in 2X SC-leu medium, pH  
6.2, with trace metals, and vitamins. After 24  
20       hours, YEP medium-containing glycerol was added to a  
final concentration of 2X YET/3% glycerol. Approxi-  
mately 24 hr later, cells were harvested, washed,  
and stored at -70°C.

**HUMAN PHOSPHODIESTERASE PREPARATIONS****Phosphodiesterase Activity Determinations**

30                   Phosphodiesterase activity of the prepara-  
tions was determined as follows. PDE assays utiliz-  
ing a charcoal separation technique were performed  
essentially as described in Loughney et al. (1996).  
In this assay, PDE activity converts [32P]cAMP or

- 83 -

[32P]cGMP to the corresponding [32P]5'-AMP or [32P]5'-GMP in proportion to the amount of PDE activity present. The [32P]5'-AMP or [32P]5'-GMP then was quantitatively converted to free [32P]phosphate and unlabeled adenosine or guanosine by the action of snake venom 5'-nucleotidase. Hence, the amount of [32P]phosphate liberated is proportional to enzyme activity. The assay was performed at 30°C in a 100  $\mu$ L reaction mixture containing (final concentrations) 40 mM Tris HCl (pH 8.0), 1  $\mu$ M ZnSO<sub>4</sub>, 5 mM MgCl<sub>2</sub>, and 0.1 mg/mL bovine serum albumin (BSA). PDE enzyme was present in quantities that yield <30% total hydrolysis of substrate (linear assay conditions). The assay was initiated by addition of substrate (1 mM [32P]cAMP or cGMP), and the mixture was incubated for 12 minutes. Seventy-five (75)  $\mu$ g of *Crotalus atrox* venom then was added, and the incubation was continued for 3 minutes (15 minutes total). The reaction was stopped by addition of 200  $\mu$ L of activated charcoal (25 mg/mL suspension in 0.1 M NaH<sub>2</sub>PO<sub>4</sub>, pH 4). After centrifugation (750 X g for 3 minutes) to sediment the charcoal, a sample of the supernatant was taken for radioactivity determination in a scintillation counter and the PDE activity was calculated.

#### Purification of PDE5 from *S. cerevisiae*

Cell pellets (29 g) were thawed on ice with an equal volume of Lysis Buffer (25 mM Tris HCl, pH 8, 5 mM MgCl<sub>2</sub>, 0.25 mM DTT, 1 mM benzamidine, and 10  $\mu$ M ZnSO<sub>4</sub>). Cells were lysed in a Microfluidizer<sup>®</sup> (Microfluidics Corp.) using nitrogen at

- 84 -

20,000 psi. The lysate was centrifuged and filtered through 0.45  $\mu$ m disposable filters. The filtrate was applied to a 150 mL column of Q SEPHAROSE<sup>®</sup> Fast-Flow (Pharmacia). The column was washed with 1.5 volumes of Buffer A (20 mM Bis-Tris Propane, pH 6.8, 1 mM MgCl<sub>2</sub>, 0.25 mM DTT, 10  $\mu$ M ZnSO<sub>4</sub>) and eluted with a step gradient of 125 mM NaCl in Buffer A followed by a linear gradient of 125-1000 mM NaCl in Buffer A. Active fractions from the linear gradient were applied to a 180 mL hydroxyapatite column in Buffer B (20 mM Bis-Tris Propane (pH 6.8), 1 mM MgCl<sub>2</sub>, 0.25 mM DTT, 10  $\mu$ M ZnSO<sub>4</sub>, and 250 mM KCl). After loading, the column was washed with 2 volumes of Buffer B and eluted with a linear gradient of 0-125 mM potassium phosphate in Buffer B. Active fractions were pooled, precipitated with 60% ammonium sulfate, and resuspended in Buffer C (20 mM Bis-Tris Propane, pH 6.8, 125 mM NaCl, 0.5 mM DTT, and 10  $\mu$ M ZnSO<sub>4</sub>). The pool was applied to a 140 mL column of SEPHACRYL<sup>®</sup> S-300 HR and eluted with Buffer C. Active fractions were diluted to 50% glycerol and stored at -20°C.

The resultant preparations were about 85% pure by SDS-PAGE. These preparations had specific activities of about 3  $\mu$ mol cGMP hydrolyzed per minute per milligram protein.

- 85 -

Inhibitory Effect on cGMP-PDE

cGMP-PDE activity of compounds of the present invention was measured using a one-step assay adapted from Wells et al., *Biochim. Biophys. Acta*, 384, 430 (1975). The reaction medium contained 50 mM Tris-HCl; pH 7.5, 5 mM magnesium acetate, 250  $\mu$ g/ml 5'-Nucleotidase, 1 mM EGTA, and 0.15  $\mu$ M 8-[H<sup>3</sup>]-cGMP. Unless otherwise indicated, the enzyme used was a human recombinant PDE5 (ICOS Corp., Bothell, Washington).

Compounds of the invention were dissolved in DMSO finally present at 2% in the assay. The incubation time was 30 minutes during which the total substrate conversion did not exceed 30%.

The IC<sub>50</sub> values for the compounds examined were determined from concentration-response curves typically using concentrations ranging from 10 nM to 10  $\mu$ M. Tests against other PDE enzymes using standard methodology showed that compounds of the invention are selective for the cGMP-specific PDE enzyme.

Biological Data

The compounds according to the present invention were typically found to exhibit an IC<sub>50</sub> value of less than 500 nM. *In vitro* test data for representative compounds of the invention is given in the following table:

- 86 -


Table 1. <i>In vitro</i> results	
Example	PDE5 IC <sub>50</sub> (nM)
1	48.1
2	401.7
3	1.2
4	6.0
5	288.0
6	151.0
7	7.0
8	37.0

Obviously, many modifications and variations of the invention as hereinbefore set forth can be made without departing from the spirit and scope thereof, and, therefore, only such limitations should be imposed as are indicated by the appended claims.

- 87 -


## WHAT IS CLAIMED IS:

1. A compound having a formula



wherein  $R^1$  is selected from the group consisting of hydrogen,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $\text{halo}C_{1-6}$ alkyl,  $C_{3-8}$ cycloalkyl, hetero $C_{3-8}$ cycloalkyl,  $C_{3-8}$ cycloalkyl $C_{1-3}$ alkyl, aryl $C_{1-3}$ alkyl, and heteroaryl $C_{1-3}$ alkyl;

$R^2$  is selected from the group consisting of an optionally substituted monocyclic aromatic ring selected from the group consisting of benzene, thiophene, furan, and pyridine, and an optionally substituted bicyclic ring



wherein the fused ring A is a 5- or 6-membered ring, saturated or partially or fully unsaturated, and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulfur, and nitrogen;



- 88 -

$R^3$  is selected from the group consisting of hydrogen and  $C_{1-6}$ alkyl,

or  $R^1$  and  $R^3$  together represent a 3- or 4-membered alkyl or alkenyl chain component of a 5- or 6-membered ring;

$R^4$ , independently, is selected from the group consisting of

aryl,

Het,

$C_{3-8}$ cycloalkyl,

$YC_{3-8}$ cycloalkyl (wherein Y is oxygen, sulfur, or  $NR^a$ ),

$C(=O)R^a$ ,

$OC(=O)R^a$ ,

$C(=O)OR^a$ ,

$C_{1-4}$ alkylene $NR^aR^b$ ,

$C_{1-4}$ alkyleneHet,

$C_{1-4}$ alkylene $C(=O)OR^a$ ,

$C(=O)NR^aSO_2R^c$ ,

$C(=O)C_{1-4}$ alkyleneHet,

$C(=O)NR^aR^b$ ,

$C(=O)NR^aR^c$ ,

$C(=O)NR^aC_{1-4}$ alkylene $OR^b$ ,

$C(=O)NR^aC_{1-4}$ alkyleneHet,

$OR^a$ ,

$OC_{1-4}$ alkylene $C(=O)OR^a$ ,

$OC_{2-4}$ alkylene $NR^aR^b$ ,

$OC_{1-4}$ alkyleneHet,

$OC_{2-4}$ alkylene $OR^a$ ,

$OC_{1-4}$ alkylene $NR^aC(=O)OR^b$ ,

$NR^aR^b$ ,

$NR^aC_{1-4}$ alkylene $NR^aR^b$ ,

$NR^aC(=O)R^b$ ,

- 89 -

$\text{NR}^a\text{C}(=\text{O})\text{NR}^a\text{R}^b$ ,  
 $\text{N}(\text{SO}_2\text{C}_{1-4}\text{alkyl})_2$ ,  
 $\text{NR}^a(\text{SO}_2\text{C}_{1-4}\text{alkyl})$ ,  
 nitro ( $\text{NO}_2$ ),  
 trifluoromethyl,  
 trifluoromethoxy,  
 cyano ( $\text{CN}$ ),  
 $\text{SO}_2\text{NR}^a\text{R}^b$ ,  
 $\text{SO}_2\text{R}^a$ ,  
 $\text{SOR}^a$ ,  
 $\text{SR}^a$ ,  
 and  $\text{OSO}_2\text{CF}_3$ ;

$\text{R}^5$  is selected from the group consisting of hydrogen, halogen, and  $\text{C}_{1-6}\text{alkyl}$ ;

or  $\text{R}^4$  and  $\text{R}^5$  are taken together with the carbon atoms to which they are attached to form a 5-, 6-, or 7-membered ring, saturated or partially or fully unsaturated, optionally substituted and optionally containing one or two heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen;

$\text{R}^a$  is selected from the group consisting of hydrogen,  $\text{C}_{1-6}\text{alkyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ , aryl,  $\text{arylC}_{1-3}\text{alkyl}$ ,  $\text{C}_{1-3}\text{alkylenearyl}$ , and Het;

$\text{R}^b$  is selected from the group consisting of hydrogen,  $\text{C}_{1-6}\text{alkyl}$ , aryl,  $\text{arylC}_{1-3}\text{alkyl}$ ,  $\text{C}_{1-3}\text{alkylenearyl}$ , and Het;

$\text{R}^c$  is phenyl or  $\text{C}_{4-6}\text{cycloalkyl}$ , either optionally substituted with one or more substituent selected from the group consisting of halo,  $\text{C}(=\text{O})\text{OR}^a$ , and  $\text{OR}^a$ ;

Het is a 5- or 6-membered heterocyclic group, saturated or partially or fully unsaturated,

- 90 -


containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with C<sub>1-4</sub>alkyl or C(=O)OR<sup>b</sup>;

q is 1, 2, or 3; and

pharmaceutically acceptable salts and hydrates thereof.

- 91 -

2. The compound of claim 1 represented by the formula



and pharmaceutically acceptable salts and solvates thereof.


3. The compound of claim 1 wherein R<sup>1</sup> is selected from the group consisting of hydrogen, C<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkyl, C<sub>3-6</sub>cycloalkyl, and C<sub>3-6</sub>cycloalkylmethyl.

4. The compound of claim 1 wherein R<sup>3</sup> is hydrogen.

5. The compound of claim 1 wherein R<sup>2</sup> is an optionally substituted bicyclic ring selected from the group consisting of naphthalene, indene, benzoxazole, benzothiazole, benzisoxazole, benzimidazole, quinoline, indole, benzothiophene, and benzofuran.



- 92 -

6. The compound of claim 1 wherein  $R^2$  is




and wherein  $n$  is an integer 1 or 2, and  $X$ , independently, are  $C(R^a)_2$ ,  $O$ ,  $S$ , or  $NR^a$ .

7. The compound of claim 1 wherein  $R^2$ , substituted or unsubstituted, is selected from the group consisting of



, and



- 93 -

8. The compound of claim 7 wherein  $R^2$  is substituted with a substituent selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $OR^a$ ,  $CO_2R^a$ , halomethyl, halomethoxy, cyano, nitro, and  $NR^aR^b$ .

9. The compound of claim 1 wherein  $R^4$  is selected from the group consisting of aryl, trifluoromethyl, trifluoromethoxy,  $C(=O)R^a$ ,  $C(=O)OR^a$ ,  $C(=O)NR^aR^b$ ,  $C(=O)NR^aR^c$ ,  $OR^a$ , CN,  $C_{1-4}$ alkylene $NR^aR^b$ ,  $OC_{2-4}$ alkylene $NR^aR^b$ ,  $SO_2NR^aR^b$ ,  $OC(=O)R^a$ ,  $NR^aR^b$ , Het,  $C_{3-8}$ -cycloalkyl, and  $C_{3-8}$ cycloalkyl.

10. The compound of claim 9 wherein  $R^4$  is selected from the group consisting of  $CH_2NR^aR^b$ , aryl, CN,  $OR^a$ ,  $C(=O)OR^a$ , and  $NR^aR^b$ .

11. The compound of claim 10 wherein  $R^4$  is selected from the group consisting of  $CH_2NH_2$ ,  $CO_2H$ ,  $CO_2CH_3$ ,  $C_6H_5$ ,  $OCH_2C_6H_5$ , OH, CN, and  $OCH_3$ .



12. The compound of claim 1 wherein  $R^4$  and  $R^5$  are taken together to form a 6-membered saturated or unsaturated ring, optionally substituted and optionally containing one or two heteroatoms.

13. The compound of claim 12 wherein  $R^4$  and  $R^5$  are taken together to form a phenyl ring.




14. The compound of claim 1 wherein  $R^5$  is hydrogen.

- 94 -

15. A compound selected from the group consisting of






- 95 -









- 96 -





- 97 -



- 98 -



- 99 -



and pharmaceutically acceptable salts and solvates thereof.

16. A pharmaceutical composition comprising a compound of claim 1, together with a pharmaceutically acceptable diluent or carrier.

- 100 -

17. A method of treating a male or female animal in the treatment of a condition where inhibition of a cGMP-specific PDE is of a therapeutic benefit comprising treating said animal with an effective amount of a pharmaceutical composition comprising a compound of claim 1, together with a pharmaceutically acceptable diluent or carrier.

18. The method of claim 17 wherein the condition is male erectile dysfunction.

19. The method of claim 18 wherein the treatment is an oral treatment.

20. The method of claim 17 wherein the condition is female sexual arousal disorder.

21. The method of claim 20 wherein the treatment is an oral treatment.

- 101 -

22. The method of claim 17 wherein the condition is selected from the group consisting of stable angina, unstable angina, variant angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, malignant hypertension, pheochromocytoma, acute respiratory distress syndrome, congestive heart failure, acute renal failure, chronic renal failure, atherosclerosis, a condition of reduced blood vessel patency, a peripheral vascular disease, a vascular disorder, thrombocythemia, an inflammatory disease, myocardial infarction, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, peptic ulcer, a gut motility disorder, postpercutaneous transluminal coronary angioplasty, carotid angioplasty, post-bypass surgery graft stenosis; osteoporosis, preterm labor, benign prostatic hypertrophy, and irritable bowel syndrome.

23. A method of treating a condition where inhibition of a cGMP-specific PDE is of therapeutic benefit, in a human or a nonhuman animal body, comprising administering to said body a therapeutically effective amount of a compound of claim 1.

24. A method for the curative or prophylactic treatment of male erectile dysfunction or female sexual arousal disorder, comprising administration of an effective dose of a compound of claim 1, and pharmaceutically acceptable salts and solvates thereof, to an animal.

- 102 -

25. Use of a compound of claim 1 for the manufacture of a medicament for the curative or prophylactic treatment of a condition where inhibition of a cGMP-specific PDE is of a therapeutic benefit.

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 December 2001 (13.12.2001)


PCT

(10) International Publication Number  
**WO 01/094345 A3**

- (51) International Patent Classification<sup>7</sup>: C07D 471/14, A61K 31/4985, A61P 15/10
- (21) International Application Number: PCT/US01/15936
- (22) International Filing Date: 15 May 2001 (15.05.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/210,137 7 June 2000 (07.06.2000) US
- (71) Applicant (for all designated States except US): LILLY ICOS LLC [US/US]; 1209 Orange Street, Wilmington, DE 19801 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): ORME, Mark, W. [US/US]; 4235 Francis Avenue, #203, Seattle, WA 98103 (US). SAWYER, Jason, Scott [US/US]; 5718 North Winthrop Avenue, Indianapolis, IN 46220 (US). DAUGAN, Alain, Claud-Marie [FR/FR]; Centre de Recherche Glaxo Smith Kline Courtaboeuf, ZA de Courtaboeuf, 25, avenue de Quebec, F-91940 Les Ulis (FR).
- (74) Agent: NAPOLI, James, J.; Marshall, Gerstein & Borun, 6300 Sears Tower, 233 South Wacker Drive, Chicago, IL 60606 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- (88) Date of publication of the international search report:  
18 July 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/094345 A3

(54) Title: DERIVATIVES OF 2,3,6,7,12,12A-HEXAHYDROPYRAZINO[1',2':1,6]PYRIDO[3,4-B]INDOLE-1,4-DIONE



(I)

(57) Abstract: Compounds of the general structural formula (I) and use of the compounds and salts and solvates thereof, as therapeutic agents.



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/15936

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/14 A61K31/4985 A61P15/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, PAJ, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 03675 A (GLAXO WELLCOME LAB SA ;DAUGAN ALAIN CLAUDE MARIE (FR)) 6 February 1997 (1997-02-06) page 1, line 3 - line 5 page 2, formula (I) page 2, line 9 - line 20 page 4, line 25 - line 28 -----	1-25

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

16 April 2002

Date of mailing of the international search report

24/04/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Hoepfner, W

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/15936

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
W0 9703675	A	06-02-1997	AU 704955 B2	13-05-1999
			AU 6419196 A	18-02-1997
			BR 9609758 A	26-01-1999
			CA 2226784 A1	06-02-1997
			CN 1195290 A	07-10-1998
			CZ 9800033 A3	13-05-1998
			WO 9703675 A1	06-02-1997
			EP 0839040 A1	06-05-1998
			HU 9900065 A2	28-05-1999
			JP 11509221 T	17-08-1999
			NO 980153 A	10-03-1998
			PL 324495 A1	25-05-1998
			SK 3998 A3	08-07-1998
			US 6140329 A	31-10-2000